



ISSN 2320-3862
JMPS 2016; 4(6): 132-137
© 2016 JMPS
Received: 20-09-2016
Accepted: 21-10-2016

Vidjinnangni Fifamè Grâce Nadège Dedehou

Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomics Sciences University of Abomey Calavi, 01 P.O. Box 526 Cotonou - Benin

Pascal Abiodoun Olounladé

(a) Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomics Sciences University of Abomey Calavi, 01 P.O. Box 526 Cotonou - Benin

(b) Pluridisciplinary Laboratory, School management and Exploitation of livestock systems National University of Agriculture, P.O. Box 95 Ketou - Benin

Dourossimi Adam Adenilé

Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomics Sciences University of Abomey Calavi, 01 P.O. Box 526 Cotonou - Benin

Géorcelin Goué Alowanou

Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomics Sciences University of Abomey Calavi, 01 P.O. Box 526 Cotonou - Benin

Erick Virgile Bertrand Azando

National Superior School of Agricultural Sciences and Techniques of Djougou, University of Parakou, P.O. Box 73 Djougou - Benin

Mawulé Sylvie Hounzangbé-Adoté

Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomics Sciences University of Abomey Calavi, 01 P.O. Box 526 Cotonou - Benin

Correspondence

Pascal Abiodoun Olounladé

Assistant Professor,
Pluridisciplinary Laboratory,
School management and
Exploitation of livestock systems,
National University of Agriculture,
P.O. Box 95 Ketou - Benin

A review on medicinal plants of *Parkia Biglobosa* (Mimosaceae -Fabaceae) and *Pterocarpus Erinaceus* (Leguminosae – Papilionoidea)

Vidjinnangni Fifamè Grâce Nadège Dedehou, Pascal Abiodoun Olounladé, Dourossimi Adam Adenilé, Géorcelin Goué Alowanou, Erick Virgile Bertrand Azando and Mawulé Sylvie Hounzangbé-Adoté

Abstract

The use of the medicinal plants constitutes a significant resource employed by communities for their daily healthcare. *Parkia biglobosa* and *Pterocarpus erinaceus* are two plants spread in West Africa and used in traditional medicine. This study was carried out in order to summarize current knowledge on medicinal uses of these plants. Data from this study showed that the plants possessed wide-reaching pharmacological actions, including anti - diarrhoea, antimicrobial, antihypertensive, anti-inflammatory, analgesic, antiplasmodial activity. All the parts of the plants are used for the care. Phytochemical investigations revealed that some compounds as alkaloids, tannins, saponins, flavonoids, steroids, phenols, glycoside and sugars are present.

Keywords: *Parkia biglobosa*, *Pterocarpus erinaceus*, ethnopharmacological, traditional medicine

1. Introduction

Plants have formed the basis of traditional medicine system [1] which has a rapidly growing economic importance [2]. Traditional medicine refers to health practices, approaches, knowledge and beliefs incorporating plant, animal and mineral-based medicines, spiritual therapies, manual techniques and exercises, applied singularly or in combination to treat or to diagnose and prevent illnesses or maintain well-being [3]. It still remains the main recourse for a large majority of people for treating health problems. Approximately 80% of the population of the world depends according to the World Health Organization on traditional medicine, mostly herbal remedies, for their primary health care needs [4]. The African continent have a long history with the use of plants and in some African countries, up to 90% of the population rely on medicinal plants as a source of drugs [5]. This situation is justified by several reasons. Official medical attention is usually based on commercial drugs that have to be purchased with money, while a traditional medical consultancy has a much lower cost, including the consumption of the medicinal plants required [6]. The absence or inaccessibility of modern healthcare services, affordability, cultural acceptance and, under certain circumstances, effectiveness than their modern counterparts has caused a large percentage of the population to rely mostly on plant based traditional medicines for their primary health care needs [7]. These factors and a growing interest in the use of natural products and folk medicine have resulted to an increase in the demand for medicinal plants.

Like in many other developing countries, people in Bénin use medicinal plants to improve their state of health [7]. Bénin flora is too rich and various and plants are used to treat population diseases [8]. *Parkia biglobosa* Benth is a perennial deciduous tree from the savannah zone of West Africa and *Pterocarpus erinaceus* is a deciduous legume tree of African savannahs and dry forests. These plants belong to the flora of Bénin. They are important multipurpose trees and are well known in many African countries. Apart from providing building materials, wood, food, fodder, weapons and other commodities, these plants are especially important as traditional medicines. In Bénin, 47% of the various identified uses of *P. biglobosa* were medicinal [9].

The present study was set up to make the synthesis of the various uses of the two plants in traditional medicine and to support the possibility to use them like anthelmintic in the domestic ruminants to improve the productivity of the animals

2. Methods

In the present review, informations regarding medicinal properties, medicinal uses, botany and biochemical properties of *Parkia biglobosa* and *Pterocarpus erinaceus* were gathered via searching books and scientific databases including: published works on journals, PubMed, Elsevier, Google scholar, Science Direct, Springer, etc.

3. Observations and Results

3.1 Botany

Parkia biglobosa (sub-family Mimosoideae and family Fabaceae) popularly known as the African locust bean tree, is perennial tree legume. The tree grows in the savannah region of West Africa up to the southern edge of the Sahel zone 13° [10]. The fruit is slightly bent, brown indehiscent pod, 30 to 40 cm long and 2 to 3 cm wide producing up to 20 seeds.

Pterocarpus erinaceus Poir. (Leguminosae, subfamily Papilionoidea) is a small to medium-sized tree 12–15 m tall

with a diameter of 1.2–1.8 m. Exceptionally tall trees reaching 35 m height have been reported [11]. The branches are light grey. The leaves are compound, imparipinnate, about 30 cm long composed of 10 to 15 alternate or sub-opposite leaflets. The young fruits are light green and turn light brown when dry.

3.2 Medicinal uses and pharmacology studies

Parkia biglobosa and *Pterocarpus erinaceus* are medicinal plants widely used in traditional African medicine particularly in Benin, Nigeria, Burkina Faso and Ivory Coast, Togo, Mali to treat a range of diseases. The use of the two plants for this purpose related especially to the bark, the leaves and the roots. The bark was the most used parts based on the high number of uses reported. The plants are mainly used against digestive disorders, wounds, hypertension and infections (table 1) and some studies are carried out to justify these uses (table 2).

Table 1: Uses of *P. biglobosa* and *P. erinaceus* in traditional medicine.

Scientific name and Family	Plant part used	Therapeutic indication	Medicinal Preparation (and administration mode)	References
<i>Parkia biglobosa</i>	Bark	Gastric and duodenal ulcer	Powder, maceration (oral)	[12]
<i>Parkia biglobosa</i>	Bark	Depurative, diuretic	Maceration (oral)	[8]
<i>Parkia biglobosa</i>	Leaves	Abscess	Grinding (local application)	[8]
<i>Parkia biglobosa</i>	Leaves	Liver diseases, hepatic deficiency	Decoction (oral, bath)	[13]
<i>Parkia biglobosa</i>	Leaves	Diabetes	Infusion (oral)	[14]
<i>Parkia biglobosa</i>	Leaves, pods, stem bark	Wounds	Powder, decoction (local application)	[15]
<i>Parkia biglobosa</i>	Root bark, trunk bark, leaves	Digestive, nervous, cutaneous and pregnancy-birth disorders, infection and infestation, poisoning	Decoction, calcinations, trituration and maceration (oral, bath, application)	[16]
<i>Parkia biglobosa</i>	Bark, fruit pulp	High blood pressure, Yellow fever, Constipation	Bark and fruit are soaked and drunk (oral)	[17]
<i>Parkia biglobosa</i>	Stem bark, leaves	Wounds, pain, fungal infection	Decoction of bark in water (bath and/or to drink) The powder is usually dissolved in water (bath and/or to drink) or is thrown into fire (inhalation)	[18]
<i>Parkia biglobosa</i>	Roots	Coal disease	Decoction in salty water (oral)	[19]
<i>Parkia biglobosa</i>	Roots, stem bark	Diarrhoea, general weakness, abdominal pains	Powder, decoction (oral)	[20]
<i>Parkia biglobosa</i>	Seeds	Whitlow	Fermentation (local application)	[21]
<i>Parkia biglobosa</i>	Grains, barks	Anthelmintic	Grains and roots are ground, the grains are mixed with rock salt, the roots are soaked in water (oral)	[22]
<i>Parkia biglobosa</i>	Stem bark	Anti-inflammatory, antibacterial	Not reported	[23]
<i>Parkia biglobosa</i>	Seeds, leaves	Hypertension	Decoction (oral)	[24,23]
<i>Parkia biglobosa</i>	Trunk bark, pods	Diarrhoea, dysentery, cholera	Maceration (oral)	[25]
<i>Parkia biglobosa</i>	Bark, leaves, fruits	Snake or venomous animal bite	Not reported	[26]
<i>Parkia biglobosa</i>	bark	Diarrhoea and venereal diseases	Not reported	[27]
<i>Parkia biglobosa</i>	Seeds	Diabetes	Consumption in sauce (oral)	[24]
<i>Parkia biglobosa</i>	Bark	Toothaches	Fumigation	[28]
<i>Parkia biglobosa</i>	Bark	Smallpox, varicella, measles, tooth decay, toothaches, gingivitis	Infusion (oral), decoction (mouth bath)	[29]
<i>Parkia biglobosa</i>	Roots	Fever, malaria	Not reported	[30]
<i>Parkia biglobosa</i>	Fruits, barks leaves	Constipation, anorexia, rickets, icterus Mumps Haemorrhoid, ascariasis Burns	Fruits flour (oral); gargle, inhalation, fumigation of bark; maceration of fresh leaves, roast the young leaves (local application)	[31]
<i>Parkia biglobosa</i>	Stem bark	Wound, antiseptic, disinfectant, cicatrizing	Infusion (oral)	[32]
<i>Parkia biglobosa</i>	Bark	Fever, malaria	Maceration in water (oral)	[33]
<i>Pterocarpus erinaceus</i>	Bark	Nonsaanga or Saangpèlega of hen in moré language (Mossi)	Maceration in water (oral)	[19]
<i>Pterocarpus erinaceus</i>	Roots, leaves, stem bark	Diarrhoea, polyuria, rectum itching, vomiting, palpitations abdominal pains, general weakness, insomnia	Powder (oral); decoction (oral)	[20]
<i>Pterocarpus erinaceus</i>	Bark, fruit	Digestive and nervous disorders	Decoction, calcination (oral, massage, scarification)	[16]
<i>Pterocarpus erinaceus</i>	Stem bark	Eyediseases	Decoction is administered orally and used as eye drops	[34]

Table 2: pharmacological properties of *P. biglobosa* and *P. erinaceus*

Botanical Name	Biological Activity	Part Tested	Bioassay Models	Results	References
<i>Parkia biglobosa</i>	Antimicrobial Activities	ethyl acetate, ethanol and water extracts of leaves	The agar cup diffusion and dilution method	The extracts exhibited a concentration dependent antibacterial, inhibiting the growth of <i>S. aureus</i> , <i>B. cereus</i> , <i>E. coli</i> , <i>P. aeruginosa</i> , <i>A. niger</i> and <i>C. utilis</i>	[35]
		methanolic extract and aqueous fractions of the leaf, stem bark and root	agar – well diffusion method	<i>S. aureus</i> , <i>B. subtilis</i> , <i>E. coli</i> <i>P. aeruginosa</i> were inhibited by the extracts and their aqueous fractions at concentrations between 2.5 – 20 mg.	[36]
<i>Parkia biglobosa</i>	Antihypertensive effect	aqueous extract of bark of trunk	Ludwig manometer principle	The extracts decreased in a dose-depend manner the rabbit blood pressure and the effect was similar to that produced by propranolol	[37]
		Fermented seeds	Clinical, biochemical and anthropometrical, analyses in two types of people (consumption or no of seeds condiment)	Significantly decreased blood pressure and heart beat were detected in the group where the condiment is highly consumed region when compared to the non-consumption group	[38]
<i>Parkia biglobosa</i>	Anthelmintic effect	Aqueous extracts of seeds and leaves	Egg hatch assay	Produced a high hatching egg inhibition against <i>Haemonchus</i> , <i>Trichostrongylus</i> , <i>Oesophagostomum</i> and <i>Bunostomum</i> species	[39]
<i>Parkia biglobosa</i>	Antimicrobial Activities	Aqueous extract of leaves	WHO method <i>in vitro</i> micro test (Mark III)	There was a dose dependent inhibition of parasitemia in the <i>in vivo</i> tests, with maximum effect at 600 mg/kg and <i>in vitro</i> a weak and concentration-dependent activity against <i>P. falciparum</i> .	[40]
<i>Parkia biglobosa</i>	Analgesic and anti-inflammatory activities.	Hexane fraction of bark	The abdominal writhing test in mice and the hot-plate method	The extract has a marked analgesic activity when evaluated with the abdominal writhing test in mice, but, like paracetamol, was ineffective with the hot-plate method.	[41]
<i>Parkia biglobosa</i>	Anti-diabetic activity	aqueous and methanolic extracts of fermented seeds	Determination of fasting plasma glucose, total cholesterol, triglyceride, high-density lipoprotein and low-density lipoprotein	Extracts exert a hypoglycaemic effect and ameliorated the loss of bodyweight usually associated with diabetes.	[42]
<i>Pterocarpus erinaceus</i>	Antimicrobial activity	n-hexane, ethyl acetate and methanol extracts of stem bark	the cupplate method on gram positive and gram negative pathogens	The methanol fraction was the only extract that showed activity against <i>C. albicans</i> , <i>S. aureus</i> , <i>E. coli</i> , <i>B. subtilis</i> and <i>P. aeruginosa</i>	[43]
		aqueous acetone extracts	agar disc diffusion, micro-well dilution and Minimum Bactericidal Concentration assay	The extract exhibited antibacterial activity against <i>S. aureus</i> and <i>B. cereus</i>	[44]
<i>Pterocarpus erinaceus</i>	Antidiarrheal activity	Methanol leaf extract	castor oil-induced diarrhea, charcoal meal transit time and castor oil-induced enteropooling	The extract caused a significant decrease in the wet faeces, decreased the distance travelled by the charcoal meal and caused a reduction in the intraluminal fluid accumulation.	[45]
<i>Pterocarpus erinaceus</i>	Antiamaril activity	ethanolic leaves extracts	<i>in vitro</i> antimalarial tests were performed by light microscopy using Giemsa-stained smears	The extract has a moderate activity against <i>Plasmodium falciparum</i>	[46]
<i>Pterocarpus erinaceus</i>	Treatment of cognitive disorders	Aqueous bark extract	MTT colorimetric assay	Extract significantly decreased a production, displaying effects similar to those of DAPT on APP processing, but may act on another inhibition site.	[47]
<i>Pterocarpus erinaceus</i>	Anti-inflammatory effect	Methanol and dichloromethane stem bark extracts	Carrageenan induced paw edema test and croton oil induced-ear edema	Methanol extract reduced the carrageenan induced –hind paw edema. The extracts reduced the ear edema at different doses.	[48]
<i>Pterocarpus erinaceus</i>	Analgesic effect	Methanol stem bark extract	Acetic acid-induced writhing test	The extract showed an important analgesic effect against writhing induced by acetic acid injection	[48]
<i>Pterocarpus erinaceus</i>	Antioxidant potential	Aqueous, acetone leaves and barks extracts	Inhibition of free radical DPPH and Iron (III) to iron (II) reduction activity (FRAP).	The extract showed an antioxidant potential. It reduced capacity DPPH radicals which is determined by the decrease absorbance at 517 nm.	[44]
<i>Pterocarpus erinaceus</i>	Haemostatic activity	Ethanolic stem bark extract	Bleeding and clotting times determination	Administration of the extract orally revealed a significant decrease in bleeding time compared to control.	[49]
<i>Pterocarpus erinaceus</i>	Antitrypanosomal activity	solvent soluble extracts of stem bark	<i>in vivo</i> screening for antitrypanosomal activity against <i>Trypanosoma brucei brucei</i>	The extracts showed only trypanostatic effects against <i>Trypanosoma brucei brucei</i> and could not clear the parasites completely.	[50]

3.3 Phytochemical components isolated from *Parkia biglobosa* and *Pterocarpus erinaceus*

The analysis of *P. biglobosa* leaves revealed the presence of saponins, tannins, flavonoids, terpenes, phenols, sterols, isoquinoline alkaloids, indole alkaloids, cardiac glycosides and reducing sugars [40,51]. However resins, volatile oil and anthraquinones were absent [40]. Ajaiyeoba (2002) indicated the presence of saponin, tannin, flavonoids and cardiac glycosides in the leaf root and stem bark. [52] showed that coumarins and anthocyanosides are present in both bark and leaf but in different concentrations. The bark and the pods

contain parkine [53].

Phytochemical screening of the stem bark showed that *P. erinaceus* contains both condensed and hydrolysable tannins, saponins, flavonoids, terpenes, carbohydrate and phenols [43, 50]. The metabolites such as sterols, triterpenes, saponosides, anthocyanidins, coumarins and reducing compounds were isolated in stem bark and coumarins, anthocyanins, flavones and reducing compounds in the leaves by [54]. The presence of some isoflavonoids as prunetin, muningin, afromosin, tectorigenin, pseudobaptigenin is revealed in heartwood of [55].

3.4 Toxicity

The use of *P. biglobosa* as herbal medicaments in African countries and the reports on the toxicity of the plant further show that the plant is non-toxic to humans [51] but the bark and the pods of the same plant contain parkine and equally have piscicidal property [56]. The extracts of raw *P. biglobosa* seed exhibited termiticide activity [57]. No toxicity was reported for *P. erinaceus* [49, 58] showed that the supplementation of *A. guyanus* with tannin rich *P. erinaceus* or sole feeding of *P. erinaceus* did not pose any threat to the health of the animals under the conditions of the experiment.

4. Discussion

The results of this review of literature show that the populations of many countries (Bénin, Burkina-Faso, Nigeria, Ivory Coast, Togo, Mali etc.) use *P. erinaceus* and *P. biglobosa* to treat several diseases. The medicinal uses of the species were the most diversified. Particularly, the digestive system diseases (diarrhoea, dysentery, abdominal pain), diseases of the cardiovascular system, wounds, pains and burns, infectious diseases (malaria, abscesses, yellow fever, measles) are reported [28, 24, 33, 20, 16]. The scientific base of some uses of these plants was established by several studies which revealed that the plants have pharmacological properties such as antimicrobial, antihypertensive, antioxidant, anti-inflammatory, analgesic, anti-diabetic, antiplasmodial. The antimicrobial activity was determined using mainly agar diffusion method. Different extracts of both plants were prepared with different solvents. These studies showed that different extracts from leaves and stem bark in different concentrations significantly inhibited the growth of *S. aureus*, *B. cereus*, *B. subtilis*, *E. coli*, *P. aeruginosa*, *A. niger*, *C. albicans* and *C. utilis* [35, 36, 43, 44]. These activities may justify the use of these plants against infections, fungal skin diseases [16, 20, 27]. The anti-inflammatory and analgesic properties were investigated for the bark of *P. biglobosa* using the abdominal writhing test in mice and the hot-plate method and for the stem bark of *P. erinaceus* using the carrageenan induced paw edema test, croton oil induced-ear edema and acetic acid-induced writhing test. The results show that *P. biglobosa* possesses a marked analgesic activity when evaluated with the abdominal writhing test in mice but like paracetamol, was ineffective with the hot-plate method, a feature suggesting a peripheral mechanism of action [41]. *P. erinaceus* was able to reduce the carrageenan induced-hind paw edema, reduced the ear edema and showed an important analgesic effect against writhing induced by acetic acid injection [48]. These properties may explain the traditional use of *P. biglobosa* against ulcers, pains, toothaches [12, 18, 28]. The antihypertensive effect was evaluated for *P. biglobosa* using Ludwig manometer principle. The extract decreased the rabbit blood pressure [37]. Significantly decreased blood pressure and heart beat were detected in the group where the condiment is highly consumed region when compared to the non-consumption group [38]. These results justify the use of this plant by traditional healers in the high blood pressure treatment [24, 23]. *P. biglobosa* (grains, bark and leaves) is reported to be effective against the worms [22, 31, 39] showed *in vitro* the anthelmintic effect of aqueous extracts of seeds and leaves on bovine nematode eggs. The extracts produced a high hatching egg inhibition against *Haemonchus*, *Trichostrongylus*, *Oesophagostomum* and *Bunostomum* species. These results supported the anthelmintic use of this plant. The antimalarial activities are recognized by the traditional healers [33, 40] indicated that with *P. biglobosa*, there was *in vivo* a dose dependent

inhibition of parasitaemia but the *in vitro* screening demonstrated a weak and concentration-dependent activity of the extract against *P. falciparum*. [46] showed that the leaves extracts have a moderate activity against *P. falciparum*. *P. biglobosa* and *P. erinaceus* are widely used in traditional medicine to treat a range of diseases. Their medicinal properties would undoubtedly be due to the various phytochemical components which they contain.

5. Conclusion

The synthesis of current knowledge on *P. biglobosa* and *P. erinaceus* shows that they are used in many countries in Africa to treat various affections like digestive disorders, hypertension, wounds, and aches. Several studies were carried out to validate some of their therapeutic uses. These plants are used especially to improve human health and very little for animal health then we can evaluate their anthelmintic properties against gastrointestinal parasites of small ruminants Djallonké.

6. Acknowledgments

This work was supported by UEMOA through the PAES project VPMAP. Mrs V.N.G. DEDEHOU is a grateful recipient of a grant provided from MECESRS of Benin. Mr. A.P. OLOUNLADE acknowledges receiving a scholarship from University of Abomey-Calavi to undergo his Post-Doc studies. The authors sincerely thank traditional practitioners from Bénin.

7. References

1. Dramane S, Witabouna KM, Kagoyire K. Evaluation of antimicrobial and free radical scavenging activities of some bioactive taxons from Côte D'ivoire. Eur. J. Sci. Res. 2010; 40:307-317.
2. Agra MF, Freitas PF, Barbosa-Filho JM. Synopsis of the plants known as medicinal and poisonous in Northeast of Brazil. Rev. Bras. Farmacogn. 2007; 17:114-140.
3. World Health Organization. Traditional Medicine, 2003. [<http://www.who.int/mediacentre/factsheet/fs134/en/print/html>]
4. Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. J Ethnobi Ethnomed. 2006; 2:43.
5. Hostettmann K, Marston A, Ndjoko K, Wolfender JL. The Potential of African Medicinal Plants as a Source of Drugs. Current Organic Chemistry. 2000; 4:973-1010.
6. Macia Manuel J, Garcia E, Vidaurre Prem J. An ethnobotanical survey of medicinal plants commercialized in the markets of La Paz and El Alto, Bolivia. Journal of ethnopharmacology. 2005; 97:337-350.
7. Ministry of Health. Document Politique Nationale de la Médecine Traditionnelle au Bénin. Cotonou, Bénin, Ministère de la Santé, 2011, 37.
8. Adjanohoun E, Adjakidje V, Ahyi MRA, Ake Assi L, Akoeigninou A, d'Almeida J. *et al.* Contribution aux études ethnobotaniques et floristiques en République populaire du Bénin. Médecine Traditionnelle et pharmacopée. Agence de Coopération Culturelle et Technique, Paris, 1989, 895.
9. Koura K, Ganglo JC, Assogbadjo AE, Agbangla C. Ethnic differences in use values and use patterns of *Parkia biglobosa* in Northern Benin. Journal of Ethnobiology and Ethnomedicine. 2011; 7:42. <http://www.ethnobiomed.com/content/7/1/42>.

10. Campbell-platt G. African locust bean (*Parkia* species) and its West African fermented food product, dawadawa. *Ecol. Food Nutr.* 1980; 9:123-132.
11. Von Maydell HJ. Arbres et arbustes du Sahel. Leurs caractéristiques et leurs utilisations. Eschnorn, GT, 1983, 531.
12. Adjanohoun E, Adjakidjè V, Ahyi MRA, Akpagana K, Chibon P, El - Hadji A. *et al.* Contribution aux études ethnobotaniques et floristiques au Togo. Agence de coopération culturelle et technique (ACCT), Paris, 1986, 671.
13. Adjanohoun E, Ake Assi L. Aux contributions des plantes médicinales *census de Côte d'Ivoire*. Centre Nat. Fleuriste. Univ. Abidjan 1979; 358. Abo KA, Fred-Jaiyesimi AA, Jaiyessimi AEA, 2008.
15. Inngierdingen K, Nergard CS, Diallo D, Mounkoro PP, Paulsen BS, Zerbo P. *et al.* Plantes médicinales et pratiques médicales au Burkina Faso. cas des Sanan. Bois et Forêts des Tropiques. 2011; 307(1):41-53.
16. Ampitan TA. Ethnobotanical survey of medicinal plants in Biu local government area of Borno state, Nigeria. *Comprehensive Journal of Herbs and Medicinal Plants.* 2013; 2(1):7-11.
17. Grønhaug TE, Glæserud S, Skogsrud M, Ballo N, Bah S, Diallo D. *et al.* Ethnopharmacological survey of six medicinal plants from Mali, West-Africa. *J Ethnobiol Ethnomed*, 2008, 4(26).
18. Tamboura H, Kaboré H, Yaméogo SM. Ethnomédecine vétérinaire et pharmacopée traditionnelle dans le plateau central du Burkina Faso : cas de la province du Passoré. *Biotechnol. Agron. Soc. Environ.* 1998 ; 2(3):181-191.
19. Tchacondo T, Karou SD, Batawila K, Agban A, Ouro-Bang'na K, Anani KT. *et al.* Herbal remedies and their adverse effects in tem tribe traditional Medicine in Togo. *Afr J Tradit Complement Altern Med.* 2011; 8(1):45-60.
20. Assogba MN. Quelques enquêtes sur la pharmacopée traditionnelle vétérinaire en République du Bénin. 13^e Conférence de la Société Ouest Africaine de Pharmacologie; Février, 1984, 23-25.
21. Djoueché CM, Azebaze AB, Dongmo AB. Investigation of Plants Used for the Ethnoveterinary Control of Gastrointestinal Parasites in Bénoué Region, Cameroon. *Tropicicultura.* 2011; 29(4):205-211.
22. Sourabie, Kinda Yaro Nikiema JB. Ethnobotanical Survey of Medicinal Plants Used By the Traditional Medical Healers in the Villages of Bérégadougou and Fabédougou (Cascades Region, Burkina Faso). *IOSR Journal of Pharmacy.* 2013; 3(7):38-45.
23. Tra Bi Fézan H, Irié GM, N'gaman KCC, Mohou CHB Ake-Assi YA Belem B, Nacoulma BMI, Gbangou R. *et al.* Use of Non Wood Forest Products by local people bordering the "Parc National Kaboré Tambi", Burkina Faso. *The Journal of Transdisciplinary Environmental Studies*, 2007, (6)1.
24. Muhammad S, Amusa NA. The Important Food Crops and Medicinal Plants of North-western Nigeria. *Research Journal of Agriculture and Biological Sciences.* 2005; 1(3):254-260.
25. Tapsoba H, Deschamps JP Thomas LV. Saotoing P, Vroumsia T, Tchobsala FN, Tchuenguem Fohouo, AM, 2011; Malgras D. Magassouba FB, Diallo A, Kouyaté M, Mara F, Bangoura O, Camara A *et al.*
28. Kayodé J, Ige OE, Adetogo TA, Igbakin APKoné WM, Atindehou Kamanzi K. Ethnobotanical inventory of medicinal plants used in traditional veterinary medicine in Northern Côte d'Ivoire (West Africa). *South African Journal of Botany.* 2008; 74:76-84.
29. Ajaiyeoba E0. Phytochemical and antibacterial properties of *Parkia biglobosa* and *Parkia bicolor* leaf extracts. *African Journal of Biomedical Research.* 2002; 5(3):125-129.
30. Udobi CE, Onaolapo JA. Phytochemical analysis and antibacterial evaluation of the leaf stem bark and root of the African locust bean (*Parkia biglobosa*). *Journal of Medicinal Plants Research.* 2009; 3(5):338-344.
31. Kassi Y, Aka KJ, Abo KJ-C, Méa A, Néné Bi SA, Ehilé EE. Effet antihypertensif d'un extrait aqueux d'écorce de tronc de *Parkia biglobosa* (Mimosaceae) sur la pression artérielle de lapin. *Sciences & Nature.* 2008; 5(2):133-143.
32. Ognatan K, Adi K, Lamboni C, Damorou JM, Aklikokou KA, Gbeassor M. *et al.* Effect of Dietary Intake of Fermented Seeds of *Parkia biglobosa* (Jacq) Benth (African Locust Bean) on Hypertension in Bogou and Goumou-kope Areas of Togo. *Tropical Journal of Pharmaceutical Research*, 2011, 10(5).
33. Soetan KO, Lasisi OT, Agboluaje AK. Comparative assessment of *in-vitro* anthelmintic effects of the aqueous extracts of the seeds and leaves of the African locust bean (*Parkia biglobosa*) on bovine nematode eggs. *Journal of Cell and Animal Biology.* 2011; 5(6):109-112.
34. Builders M, Wannang N, Aguiyi J. Antiplasmodial activities of *Parkia biglobosa* leaves. *In vivo* and *In vitro* studies. *Annals of Biological Research.* 2011; 2(4):8-20.
35. Kouadio F, Kanko C, Juge M, Grimaud N, Jean A, N'Guessan YT. *et al.* Analgesic and anti-inflammatory activities of an extract from *Parkia biglobosa* used in traditional medicine in the Ivory Coast, *Phytother Res.* 2000; 14(8):635-637.
36. Odetola A, Akinloye O, Egunjobi C, Adekunle W, Ayoola A. Possible antidiabetic and antihyperlipidaemic effect of fermented *Parkia biglobosa* (Jacq) extract in alloxan-induced diabetic rats. *Clinical and Experimental Pharmacology and Physiology.* 2006; 33(9):808-812.
37. Gabriel AF, Onigbanjo HO. Phytochemical and Antimicrobial Screening of the Stem Bark Extracts of *Pterocarpus erinaceus* (Poir). *Nigerian Journal of Basic and Applied Science.* 2010; 18(1):1-5.
38. Konaté K, Kiendrébéogo M, Ouattara MB, Souza A, Lamien-Meda A, Nongasida Y. *et al.* Antibacterial Potential of Aqueous Acetone Extracts From Five Medicinal Plants used Traditionally to Treat Infectious Diseases in Burkina Faso. *Current Research Journal of Biological Sciences.* 2011; 3(5):435-442.
39. Ezeja MI, Ezeigbo II, Madubuike KG, Udeh NE, Ukwani IA, Akomas SC. Antidiarrheal activity of *Pterocarpus erinaceus* methanol leaf extract in experimentally-induced diarrhea. *Asian Pacific Journal of Tropical Medicine.* 2012; (2012):147-150.
40. Karou D, Dicko MH, Sanon S, Simpore J, Traore AS. Antimalarial activity of *Sida acuta* Burm. f. (Malvaceae) and *Pterocarpus erinaceus* Poir. (Fabaceae). *Journal of Ethnopharmacology.* 2003; 89:291-294.
41. Hage S, Kienlen-Campard P, Octave JN, Quetin-Leclercq J. *In vitro* screening on amyloid peptide production of plants used in traditional medicine for cognitive disorders. *Journal of Ethnopharmacology.* 2010; 131:585-591.
42. Ouédraogo N, Sawadogo RW, Tibiri A, Bayet C, Lompo M, Hay AE. *et al.* Pharmacological properties and related constituents of stem bark of *Pterocarpus erinaceus* Poir.

- (Fabaceae). Asian Pacific Journal of Tropical Medicine. 2011; 412-420.
43. Salawu OA, Aliyu M, Tijani AY. Haematological studies on the ethanolic stem bark extract of *Pterocarpus erinaceus* poir (fabaceae). African Journal of Biotechnology, 2008; 7(9):1212-1215.
 44. Mann A, Ogbadoyi EO. Evaluation of Medicinal Plants from Nupeland for Their *in vivo* Antitrypanosomal Activity. American Journal of Biochemistry. 2012; 2(1):1-6.
 45. Abioye EO, Akinpelu DA, Aiyegoro O, Adegboye MF, Oni MO, Okoh AI. Preliminary Phytochemical Screening and Antibacterial Properties of Crude Stem Bark Extracts and Fractions of *Parkia biglobosa* (Jacq.). Molecules. 2013; 18(7):84-85.
 46. Millogo-Kone H, Guissou IP, Nacoulma O, Traore AS. Comparative Study of Leaf and Stem Bark Extracts of *Parkia Biglobosa* Against Enterobacteria. Afr J Tradit Complement Altern Med. 2008; 5(3):238-243.
 47. Fafioye OO. Plants with Piscicidal Activities in Southwestern Nigeria. Journal of Fisheries and Aquatic Sciences. 2005; 5:91-97.
 48. Millogo-Kone H, Guissou IP, Nacoulma O, Traore AS. Study of the antibacterial activity of the stem bark and leaf extracts of *Parkia biglobosa* (jacq.) Benth. on *Staphylococcus aureus*. African Journal of Traditional, Complementary and Alternative Medicines. 2006; 3(2):74-78.
 49. Bevan CWL, Ekong DEU, Obasi ME. West African timbers. XIII. Extracts from the heartwood of *Amphimas pterocarpoides* and *Pterocarpus erinaceus*. J Chem SO Org. 1966; 5:509-510.
 50. Abalaka SE, Auta J. Toxic Effects of Aqueous and Ethanol Extracts of *Parkia biglobosa* Pods on *Clarias gariepinus* Adults. World Journal of Biological Research. 2010; 003(1):9-17.
 51. Femi-Ola TO, Ajibade VA, Afolabi A. Chemical composition and termicidal properties of *Parkia biglobosa* (Jacq) Benth. Journal of Biological Sciences. 2008; 8(2):494-497.
 52. Olafadehan OA. Changes in haematological and biochemical diagnostic parameters of Red Sokoto goats fed tannin-rich *Pterocarpus erinaceus* forage diets. Vet. Arhiv. 2011; 81:471-483.