



ISSN (E): 2320-3862
ISSN (P): 2394-0530
NAAS Rating: 3.53
JMPS 2019; 7(2): 30-34
© 2019 JMPS
Received: 16-01-2019
Accepted: 20-02-2019

Nur Akter Habiba
Physical Pharmacy and
Pharmaceutical Analysis Lab,
Department of Pharmacy,
BRAC University, Dhaka,
Bangladesh

Nahid Akter
Research Lab, Department of
Pharmacy, BRAC University,
Dhaka, Bangladesh

Mahia Ferdushi
Department of Pharmacy, East
West University, Dhaka,
Bangladesh

Tanjina Afrin
Department of Pharmacy, East
West University, Dhaka,
Bangladesh

Monika Nasrin Munni
Department of Pharmacy,
BRAC University, Dhaka,
Bangladesh

Maksuda Akter
Department of Pharmacy,
BRAC University, Dhaka,
Bangladesh

Correspondence
Nur Akter Habiba
Physical Pharmacy and
Pharmaceutical Analysis Lab,
Department of Pharmacy,
BRAC University, Dhaka,
Bangladesh

Investigation of Anthelmintic and Insecticidal activity of *Goniothalamus sesquipedalis* plant in different extracts

Nur Akter Habiba, Nahid Akter, Mahia Ferdushi, Tanjina Afrin, Monika Nasrin Munni and Maksuda Akter

Abstract

Within different extract of *Goniothalamus sesquipedalis* the evaluation of anthelmintic and insecticidal activity conducted in present studies. For anthelmintic activity, selected animal was *Pheretima posthuma* for testing while 25, 50, 100 mg/ml concentrations of specimen were tested in the bioassay, from which time of paralysis and time of death of worms were estimated. Assay of insecticidal action was executed against *Sitophilus oryzae* to assess the mortality rate. MEE, EE, ACE, CHE, PEE and NHE extract were there for the activity determination. In anthelmintic study, extract exhibition and activity in dose-dependent manner showing higher the concentration, faster the effect. Extract of EE, PEE, CHE and MEE in case of 100 mg/ml concentration showed its paralytic effect followed by death within a short period of time among which EE extract gave the best result which only took approx. 22.33 minutes to show paralytic effect and 33 minutes for death occurrence. In insecticidal activity, all extracts showed efficacious activity with 100% mortality of rice insects *Sitophilus oryzae* at 80 mg/ml concentration in 24hrs, proving the activity also followed dose dependent pattern. 80% mortality rate observed in case of 80 mg/ml concentration of ACE extract in 12 hr whereas MEE, CHE and NHE extracts showed 60 % and PEE showed 70% mortality rate in treated rice insects. The present study shows that all the extracts of *Goniothalamus sesquipedalis* are found to retain anthelmintic and insecticidal activities.

Keywords: *Goniothalamus sesquipedalis*, anthelmintic, insecticidal, *Pheretima posthuma*, *Sitophilus oryzae*

Introduction

Plants are considered as wellspring of solution for a huge number of years. In various ailments plants and phyto-item keep an essential job in the treatment. Medication disclosure from plants is finished by different strategies like; herbal, ethnobotanicals, phytochemical and natural and concoction detachment which are multi-disciplinary methodology. Be that as it may, in spite of these perceptions, approximately 80% of the world's inhabitants' issues are treated by restorative home grown sedate for their essential wellbeing care [1]. These days the events of parasitic diseases are expanding widely. In spite of having various anti-infection agents accessible to experience diseases, anti-infection agent obstruction has constrained its utilization. Accordingly an elective alternative can be phytotherapy to beat nematodes. In the meantime, to attack distinctive pathogenic organism's therapeutic plants can be a potential decision to vanquish irresistible infections [2]. This plant *Goniothalamus sesquipedalis* lies under the family Annonaceae. In this way, the extracts of *Goniothalamus sesquipedalis* was utilized to decide the insecticidal and anthelmintic property of the plant.

Materials and methods

Collection of the plant: The plant *Goniothalamus sesquipedalis* was collected from the local area of Chottogram, Bangladesh in the month of June, 2017. The plant was freed from materials like dust, dirt, polen. Then the plant was identified by Scientific Officer, Bangladesh National Herbarium, Mirpur, Dhaka and a voucher specimen has been deposited (DACB:42,930) for further reference.

Extraction of plant material: The plant parts were dried under sun for a few days and finally oven dried to remove all the moisture content. Then the plant parts were crushed to coarse consistency. The coarse plant parts were extracted in a decreasing polarity order.

The coarse plant material (900g) was taken and soaked with 1500 ml of methanol for 3 consecutive days at 25°C. The extract was filtered and the filtrate was kept for further extraction. In the same manner the residue was soaked in different solvents by polarity decreasing order.

Methanol > Ethanol > Acetone > Chloroform > Petroleum ether > n-hexane

For every case, the extract was preserved and solvent evaporation was done by using rotary evaporator. Finally, all the extracts of *Goniothalamus sesquipedalis* was kept under laminar airflow for protecting it from any type of contamination.

Drugs and chemicals: Albendazole was assorted from Eskayef Bangladesh Ltd.; 0.9% sodium chloride solution (Normal saline) was purchased from IBN SINA Pharmaceutical Industry Ltd., Bangladesh and other reagents were of analytical grade.

Earthworms and Insects: *Pheretima posthuma* (*P. posthuma*), a species of adult earthworm has pretty high level of semblance with the human intestinal round worm parasite from anatomical and physiological point of view. Which lead to choose this particular earth worm for the evaluation of *in vitro* anthelmintic activity. Mohakhali area of Dhaka, Bangladesh were chosen for the collection of the adult earthworms and moist soil was preferred as collection site. All kinds of fecal matters were removed using saline water. The earthworms were of 3-5 cm in length and 0.1-0.2 cm in width as per in size. A very known and communal rice weevil, *Sitophilus oryzae* (*S. oryzae*), was collected from a local rice godown Bangladesh for detecting the insecticidal activity of *Goniothalamus sesquipedalis*. For the insect collection, old rice were preferred.

Anthelmintic activity: Development of anthelmintic resistance in helminthes reported in a number of countries gives a clear indication that control programs based exclusively on their use are not sustainable [11-15]. The development of integrated programs to control helminths is vital, but such control programs require viable alternatives to the use of anthelmintics [16]. *Pheretima posthuma*, an adult earthworm was chosen for the dtermination of anthelmintic activity. For every particular extract of *Goniothalamus sesquipedalis* sample solution was prepared at 25, 50 and 100mg/ml concentration in normal saline water and for each concentration approximately six similar size of earthworms (*Pheretima posthuma*) were placed in each beaker containing 50ml of above test solutions of extract. As a reference standard, Albendazole (10 mg/ml) was used and normal saline water as control. Solutions of each concentration was freshly prepared just before conduction the experiment. The time required for the paralysis was noted when there was no movement from the worms despite these being shaken vigorously [17]. Time for death of worms were recorded after ascertaining that worms neither moved when shaken vigorously nor when dipped in warm water (50°C).

Insecticidal activity: Insecticidal activity of many plants against several insect pests has been demonstrated [18-20]. Yang and Tang (1988) reviewed the plants used for pest insect control and found that there is a strong connection between medicinal and pesticidal plants [21]. For beginning the surface film activity test petri dishes of 60 mm were taken for the extract and their replication. For every particular extract of

Goniothalamus sesquipedalis sample solution was prepared at (2.5, 5, 10, 20, 40, 50, 60, 70, 80 mg/ml). Now in every respective petri dish the prepared sample solutions were poured into the lower part of it and kept for a few moments for it to dry out. Then insects were released in each of the treated petri dish. A control petri dish was also kept in the similar manner. Soon performing all the steps mentioned above, the petri dishes were kept in a sealed environment for the observance of the insecticidal activity. The mortality was observed first at 30 minutes and then after 12 and 24 hours of exposure. After 24 hours the data was recorded. For the verification of the movement and death of insects, a simple microscope was used. In some cases hot needle was taken closer to the insect bodies (without movement) for the confirmation of death. Recovery of the insects if occurred was also taken into consideration. The mortality rates of *Sitophilus oryzae* adults were corrected by the Abbott's formula [22].

$$\% \text{ of Mortality} = \left[\frac{Ca - Ta}{Ca} \right] \times 100$$

Where,

Ca = No. of live control insects after treatment

Ta = No. of live test insects after treatment

Statistical Analysis: All assays were performed in triplicate under strict aseptic conditions to ensure consistency of all findings. For each extract triplicate data was taken and the final data was taken by the triplicate data's mean \pm SD (Standard Deviation), which was analyzed by Microsoft excel. The bar diagram was designed using Prism software.

Results and discussion

Anthelmintic activity: Although there has been a remarkable advancement in the field of human medicines observed, still several infectious diseases triggered by viruses, bacteria, fungi, parasites, insects, rodents and helminthes are becoming a threat to communities' well-being. To be more specific developing countries are at higher risk for their persistence. Exclusively, children and teenagers are susceptible to infectious diseases. The comparative anthelmintic drug resistance incidence has led to the increasing demand of the natural anthelmintic as potential substitute for diseases management [23, 24]. The present study demonstrates the anthelmintic activity of the plant extract of *Goniothalamus sesquipedalis* using adult earthworms because of high level of semblance with the human intestinal round worm parasite. When the earthworms (*Pheretima posthuma*) were exposed to the *Goniothalamus sesquipedalis* extract, they started gradually losing their motility by providing stimulant effect primarily.

Ultimately, dose-dependent paralysis was observed followed by occurrence of death in each crude extract containing 25, 50 and 100mg/ml. Table 1 represents the effects of *Goniothalamus sesquipedalis* in different extracts. The extract denoted anthelmintic activity in dose-dependent manner describing higher concentration of each extract exhibited paralytic effect much earlier and resulted in death within a short period of time. As per the data denoted in Table 1, it reflects that at 25mg/ml concentration in all extracts (MEE, EE, ACE, CHE, PEE and NHE) the time for paralysis and ultimately the time required for the death worms took more time comparing to 50 and 100mg/ml concentration. The standard reference, albendazole showed strong anthelmintic action. As a whole, different extracts of *Goniothalamus sesquipedalis* showed anthelmintic activity in a dose-

dependent manner [24]. Previous phytochemical screening reported extract in different extracts of *Goniothalamus sesquipedalis* contain carbohydrates, tannins, flavonoids, proteins, alkaloids and [25]. Tannins being polyphenolic compounds are known to have anthelmintic activities. The reports on the anthelmintic effect of tannins is due to binding of tannins to free proteins in the gastrointestinal tract of host

animal or glycoprotein on the cuticle of the parasite and may be responsible for death [26]. Secondary metabolites like flavonoids also possesses anthelmintic activity but their exact mechanism is yet to be known [27]. So the presence of tannins and secondary metabolites like flavonoids presence in the extract of *Goniothalamus sesquipedalis* produced similar anthelmintic effects.

Table 1: Effect of different extracts of *Goniothalamus sesquipedalis* on *Pheretima posthuma*

Group	Dose (mg/ml)	No. of worms	Time taken for paralysis	Time taken for death
Control	20ml	6	---	---
Standard	10	6	28.33±1.15	35±1.73
MEE	100	6	27±2.64	43.33±2.52
	50	6	32.66±1.52	55±2.65
	25	6	51.33±1.52	72.66±1.15
EE	100	6	22.33±2.08	33±2
	50	6	43±2.64	63.66±3.51
	25	6	46.66±3.21	75.66±4.50
ACE	100	6	31±2	46.33±2.30
	50	6	39.33±1.52	52.33±3.05
	25	6	50.66±2.51	70.66±2.52
CHE	100	6	23.66±4.72	32±2
	50	6	35.66±3.05	47.33±1.53
	25	6	44.33±1.15	55.33±2.52
PEE	100	6	31.33±2.08	40.33±1.53
	50	6	37.66±1.15	55.66±2.52
	25	6	44±1	62.33±1.15
NHE	100	6	23.66±3.05	34.33±2.52
	50	6	34.66±4.04	47.66±1.53
	25	6	47.66±0.57	62±2.65

Control group received saline water 20 ml, standard group received Albendazole 10 mg/ml and test groups MEE, EE, ACE, CHE, PEE and NHE were treated with 25, 50 and 100 mg/ml of the extracts respectively. MEE= Methanol extract,

EE= Ethanol extract ACE = Acetone extract, CHE= Chloroform extract, PEE = Petroleum ether extract, NHE= n-hexane extract.




Fig 1: Graphical presentation of time vs. occurrence of paralysis




Fig 2: Graphical presentation of time vs. occurrence of death

Insecticidal activity: A major interference in the use of insecticides are resistance developed by insects to fight against diseases due to its fewer efficacies. So a potent substitution can be natural insecticides prepared from plants since they contain several bioactive chemicals [28]. The present study demonstrates the insecticidal activity of the plant extract of *Goniothalamus sesquipedalis* against rice induced insect named *Sitophilus oryzae*. For every particular extract of *Goniothalamus sesquipedalis* sample solution was prepared at (2.5, 5, 10, 20, 40, 50, 60, 70, 80 mg/ml) and for evaluation was observed for 12 and 24hr. Among all the extracts, ACE

showed the highest mortality (80%) whereas MEE, CHE, NHE extracts provided % percentage of mortality of (60%) and PEE (70%) at 80mg/ml in 12 hours (Table 3). After 24 hours, at 80 mg/ml in all extracts exhibited 100% (Table 3). The mortality of the insect was found to be concentration dependent. It is observed that the carbohydrates, saponins, phytosterols, phenols, flavonoids and tannins are having mosquito larvicidal activity [29]. So the presence of tannins and secondary metabolites like flavonoids presence in the extract of *Goniothalamus sesquipedalis* plant may describe the toxic effects on the insects.

Table 2: Insecticidal activity of *Goniothalamus sesquipedalis* On *S. Oryzae* (Death of Insects)

Sample Conc mg/ml	No. of insect taken	No. of Insects Dead											
		MEE		EE		ACE		CHE		PEE		NHE	
		12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs
2.5	10	0	1	0	0	0	1	1	3	0	2	0	0
5	10	1	1	1	0	0	2	1	3	1	2	0	2
10	10	1	3	1	3	2	4	2	4	2	3	1	3
20	10	2	4	2	3	3	5	3	4	2	4	3	4
40	10	2	5	2	6	5	6	4	6	3	5	3	4
50	10	3	5	3	6	5	9	6	7	5	6	3	6
60	10	4	8	3	7	9	10	6	8	6	7	5	7
70	10	5	7	5	10	10	10	7	8	6	8	5	10
80	10	6	10	5	10	10	10	7	10	7	10	6	10

Table 3: Insecticidal activity of *Goniothalamus sesquipedalis* On *S. Oryzae* (% of Mortality)

Sample Conc mg/ml	% of Mortality											
	MEE		EE		ACE		CHE		PEE		NHE	
	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs	12 hrs	24 hrs
2.5	0	10	0	0	0	10	10	30	0	20	0	0
5	10	10	10	0	0	20	10	30	10	20	0	20
10	10	30	10	30	20	40	20	40	20	30	10	30
20	20	40	20	30	30	50	30	40	20	40	30	40
40	20	50	20	60	50	60	40	60	30	50	30	40
50	30	50	30	60	50	90	60	70	50	60	30	60
60	40	70	30	70	90	100	60	80	60	70	50	70
70	50	80	50	100	100	100	70	80	60	80	50	100
80	60	100	50	100	80	100	70	100	70	100	60	100

Conclusion

The present study indicates the different plant extract of *Goniothalamus sesquipedalis* possess varying degree of anthelmintic activity and toxic principles with presence of insecticidal activity. The anthelmintic study shows, the acetone extract provides the fastest paralysis effect (22.33 minutes) followed by the fastest death time (33 minutes) in 100 mg/ml concentration. Whereas chloroform and n-hexane extract provides the slowest paralysis time and death occurrence time 100 mg/ml concentration. The insecticidal study demonstrates that the acetone extract provides the highest % percentage of mortality of 80% in 12 hours at 80 mg/ml concentration comparing with the other extract. Whereas MEE, CHE, NHE extracts provided % percentage of mortality of 60% and PEE 70% in 12 hours at 80 mg/ml concentration. All the extract provided 100% mortality in 24 hours at 80 mg/ml concentration. Therefore, the observed biological activities can play a great role for further detailed investigation of other particular activities of different plant extracts to signify the research on medicinal plants.

References

1. Ali MR, Hossain M, Runa JF, Hasanuzzaman M. Preliminary cytotoxic activity of different extracts of *Averrhoa bilimbi* (fruits). *International Current Pharmaceutical Journal*. 2013; 2(3):83-84.
2. Dash PR, Mou KM, Erina IN, Ripa FA, Masud KNA, Ali MS. Study of anthelmintic and insecticidal activities of different extracts of *Kaempferia galanga*. *International Journal of Pharmaceutical Science and Research*. 2017; 8(2).
3. Khan B. *Encyclopedia of Flora and Fauna of Bangladesh*. Asia Society of Bangladesh, Dhaka, Edition 1. 2008; 7:392-3.
4. Hoang SV, Pieter B, Keler PJA. Uses and conservation of plant species in a national park- a case study of Ben En, Vietnam. *Eco Botany*. 2008; 62(4):574-93.
5. Lin YF, Yi Z, Zhao YH. *Chinese Dai Medicine Colorful*

Illustrations. Yunnan Nationality Press, First Edition, 2003.

6. Yang XW, Wang JS, Ma YL, Xiao HT, Zuo YQ, Lin H *et al*. Bioactive Phenols from the Leaves of *Baccaurea ramiflora*. *Planta Medica*. 2007; 73:1415-1417.
7. Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH, Rahman S. DPPH free radical scavenging activity of some Bangladeshi medicinal plants. *Journal of Medicinal Plants Research*. 2009; 3(11):875- 879.
8. Howlader A, Sultana S, Ghosh CK, Bachar SC. Oral Hypoglycemic Effect of *Baccaurea ramiflora* L. (Phyllanthaceae) Bark Methanolic Extract on Mice. *Journal of Southeast University Science and Engineering*. 2009; 3:89-92.
9. Obayed Ullah M, Urmi KF, Howlader MA, Hossain MK, Ahmed MT, Hamid K. Hypoglycemic, hypolipidemic and antioxidant effects of leaves methanolic Extract of *Baccaurea ramiflora*. *Int. J Pharm Pharm. Sci*. 2012; 4(3):266-269.
10. Usha T, Middha SK, Bhattacharya M, Lokesh P, Goyal AK. Rosmarinic Acid, a New Polyphenol from *Baccaurea ramiflora* Lour. Leaf: A Probable Compound for Its Anti-Inflammatory Activity. *Segura-Carretero A, Arráez-Román D, eds. Antioxidants*. 2014; 3(4):830-842.
11. Jackson F. Anthelmintic resistance - the state of play. *British Vet. J*. 1993; 149:123-38.
12. Sanyal PK. Gastrointestinal parasites and small ruminant production in India. In: Le Jambre, L.F. and M.R. Knox (Eds), *Sustainable Parasite Control in Small Ruminants*. Proc. Workshop Bogor, Indonesia, ACIAR Proceedings No. 1996; 74(22-25):109-112.
13. Rolfe PF. Anthelmintic resistance in Australia, its development and management. *Proc. 4th Intl. Cong. Sheep Vet., Armidale, Australia, 1997*, 51-8.
14. Van Wyk JA, Malan FS, Randles JL. How long before resistance makes it impossible to control some field strains of *Haemonchus contortus* in South Africa with

- any of the modern anthelmintics? *Vet. Parasitol.* 1997; 70:111-22.
15. Waller PJ. The global perspective of anthelmintic resistance in nematode parasites of sheep-excluding Australasia. *Proc. 4th Intl. Cong. Sheep Vet., Armidale, Australia, February. 1997, 59-63.*
 16. Waller PJ: International approaches to the concept of integrated control of nematode parasites of livestock. *Int. J Parasitol.* 1999; 29:155-64.
 17. Kumanan R, Sridhar C, Jayaveera KN, Sudha S, Rubesh Kumar S, Duganath N. Comparative Study of Anthelmintic Activity of Different Leaf Extracts of *Tecoma stans* (L.) On Adult Indian Earthworms. *International Journal of Pharmaceutical and Clinical Research.* 2015; 2(2):63-65.
 18. Jilani G, Su HCF. Laboratory studies on several plant materials as insect repellent for protection of cereal grains. *J Econ. Entomol.* 1983; 76:154-157.
 19. Isman MB. Plant essential oils for pest diseases management. *Crop Prot.* 2000; 19:603-608.
 20. Carlini CR, Grossi-de Sá MF. Plant toxic proteins with insecticidal properties: A review on their potentialities as bioinsecticides. *Toxicon.* 2002; 40:1515-1539.
 21. Yang RZ, Tangs CS. Plants used for pest control in China: a literature review. *Econ. Bot.* 1988; 42:376-406.
 22. Khatun MH, Nesa ML, Hosen AH, Bashar S, Sarker S, Islam MR. Evaluation of the antibacterial, cytotoxic and insecticidal activities of *Hibiscus sabderiffa* barks. *International journal of Pharmacy and Research.* 2015; 5(3):170-175.
 23. Kumar MBS, Kumar MCR, Bharath AC, Kumar HRV, Kekuda TRP, Nandini KC *et al.* Screening of selected biological activities of *Artocarpus lakoocha* roxb (Moraceae) fruit pericarp. *Journal of Basic and Clinical Pharmacy.* 2010; 1(4):239-245.
 24. Riaz M, Rahman N, Haq MZU. Anthelmintic and Insecticidal Activity of *Verbascum Thapsus* L. *Pakistan Journal of Zoology.* 2013; 45(6):1593-1598.
 25. Goyal AK, Mishra T, Sen A. Antioxidant profiling of Latkan (*Baccaurea ramiflora* Lour.) wine. *Indian Journal of Biotechnology.* 2013; 12(1):137-139.
 26. Mulla WA, Thorat VS, Patil RV, Burade KB. Anthelmintic activity of leaves of *Alocasia indica* Linn. *International Journal of Pharm Tech Research.* 2010; 2(1):26-30.
 27. Kerboeuf D, Riou M, Guegnard F. Flavonoids and related compounds in parasitic disease control. *Bentham Science Publisher.* 2008; 8:116-128.
 28. Kiran R, Kekuda TRP, Kumar HGP, Hosetti BB, Krishnaswamy K. Biological activities of *Sarcanthus pauciflorus*. *Journal of Applied Pharmaceutical Science.* 2013; 3(7):105-110.
 29. Khanna VG, Kannabiran K. Larvicidal effect of *Hemidesmus indicus*, *Gymnema sylvestre* and *Elipta prostrata* against *Cules quinquefasciatus* mosquito larvae. *African Journal of Biotechnology.* 2007; 6(3):307-311.