Journal of Medicinal Plants Studies
www.PlantsJournal.com

ISSN 2320-3862
JMPS 2015; 3(3): 47-52

Received: 18-03-2015
Accepted: 21-04-2015

Ashalatha
Assistant Professor,
Department of Botany, BMS College for Women, Bangalore, Karnataka, India.

Correspondence:
Ashalatha
Assistant Professor,
Department of Botany, BMS College for Women, Bangalore, Karnataka, India

Highlight how the balance and interaction between different hormones regulate plant growth and development

Ashalatha

Abstract
Growth and development in plants are regulated by a linear sequence of hormones which act as chemical signals that influence the cellular processes and reactions to the environment. Plant hormones such as auxins, gibberellins, cytokinins, ethylene and abscisic acid work in cooperation and in opposition to control various activities like cell enlargement, cell division, differentiation and maturation. Auxins used mostly are the ones that affect the elongation of cells and apical dominance with cytokines for shoot formation that affects cell division.
GAs are essential for seed germination and stem growth the activity of which can suppress the inhibitory influence of ABA controlling stress reactions and seed dormancy. Ethylene is involved in fruit maturation and probably in mechanical stimulation, it being associated with auxins' actions that control the shedding of leaves and the fading of flowers. Growth and development of plant cells imply a certain dynamic equilibrium between these hormones in order to meet the environmental challenges. This paper presents an overview of the current awareness on hormonal interactions in plants, continued research development on molecular level and its implication to agriculture.

Keywords: Plant hormones, Auxins, Cytokinins, Gibberellins, Abscisic acid (ABA), Ethylene, Growth and development, hormonal interactions, stress response, molecular mechanisms.

Introduction
Plants' growth and development involve complex interaction of hormones that are of paramount importance in creating efficient response to the changes in the environment and in achieving reproductive goals. Plant hormones, also known phytohormones are the organic compounds that affect various processes in an organism, at trace levels. The major plant hormones include auxins, gibberellins, cytokinins, ethylene and abscisic acid which have individual and overlapping functions in certain developmental processes of the plant's life span.
Some of the general functions of auxins include the following; cell elongation, apical dominance and root initiation. They assist in cell growth because they help to make the cell wall more permeable for growth, through the breakdown of bonds. They occur in at least five classes and play a major role in stem elongation, seed germination, and flowering and their actions are usually antagonistic to abscisic acid, which brings about dormancy during unfavorable conditions such as drought [2]. Cytokinins affect cell division and differentiation of shoots in contrast to inhibit apparency of shoots and regulate nutrient movement as auxins. Ethylene is a gaseous hormone that plays a significant role in fruit maturation, leaf shedding, and horns' responses to stress, often in combination with other hormones that maintain growth under unfavorable conditions.
There seems to be an interactive and distinctive relation between these hormones based on circumstances. For example, while auxins and cytokinins have cooperative effects on cell division and differentiation, Gibberellins and abscisic acid can inhibit or promote seed germination depending on the level of the other. The relatively simple substances such as auxins and cytokinins can interact with ethylene as seen in such processes as leaf senescence and abscission. New information regarding the molecular mechanisms of hormone signaling, transport, and receptor function in mammalian and non-malian species has arisen from developments in molecular biology and biotechnology [1]. Knowledge of these mechanisms might have potentials for applications in agriculture including increasing yield, stress tolerance and regulation of growth in plants.

This paper places its focus on the relationship between different plant hormones and their involved processes in the context of growth and development with special reference to the pushing off agricultural production.

The control of growth and development of plants by hormones has been among the most investigated aspects of plant biology for a long time. These developments have immensely enriched sensibility defining the machineries regulating hormonal interactions in molecular biology and genomics [3].

Related Works

Among the phytohormones regulating plant growth, auxins and, specifically, indole-3-acetic acid (IAA) have acquired the central status. People have come up with different theories to explain biosynthesis of auxin and their uses in cell enlargement, root formation, and photo climacteric processes as well as in gravitropism. Further works showed that auxin transporters e. g. PIN-FORMED (PIN) proteins are used deeply in maintaining the gradients of auxin, which in turn are important for directional growth responses. However, there are general complexities involving the signaling of auxin and interconnection of the hormone with other hormones which is still a major issue. For example, how the auxin influences the cytokinins on the control of apical dominance, a process by which the stem discourages lateral buds growth from emerging shows a clear case of how these hormones compete to control the same process.

Cytokinins with their requirement in cell division and differentiation have been given much attention. Recent findings on cytokinin perception and signal transduction are focused on the part of histidine kinase receptors as signallers of cytokinins [4]. Subsequent studies have elaborated how cytokinins influence root development and shoot emergence by a sharp (Ã) responsiveness to auxins. The antagonism between cytokinins and auxins was further elaborated where cytokines stimulate the growth of shoots and reduce root growth, which is important in normal growth reactions.

Gibberellins (GAs) are another significant group of hormones affecting stem elongation/germination of seeds / flowering. Reports on a century of gibberellin research shed the lights on the biosynthesis and the action of GA in terms of the biochemical and genetic aspects [5]. Innovative works have identified the DELLA proteins to as negative components of GA signaling and impact growth responses with the involvement of other hormonal pathways [6]. This interaction is especially observed in stress circumstances under which GA levels are regulated in an effort to maintain growth as well as survival.

The major function of abscisic acid is linked with stress conditions, especially in the case of drought and salinity. Reviews of ABA signaling networks demonstrate the major components concerned in perceiving and responding to ABA [7]. ABA involvement in stomatal closure and seed dormancy regulation was discussed in detail, whereas the most recent data regarding molecular bases of these processes was presented. Ethylene and gibberellins interact with ABA in order to regulate growth and stress response mechanisms [8].

Ethylene is a gaseous plant hormone discussing various processes having relation with fruit ripening, leaf senescence, and stress responses. The several steps advancing our understanding of the essential focus that ethylene plays in plant development explain the signaling pathways and molecular processes. Thus, the effect of ethylene on different processes is regulated by its interaction or antagonism with other hormones including ABA and cytokinins [10]. For example, the hormone ethylene works together with ABA to control the process of stomatal closure under the conditions of drought stress, which demonstrates the cooperation of these hormones in the processes of stress response.

These hormones are related in action and function in a feedback system that constantly regulates the plant's reactions to the stimuli in the environment. New investigations have used molecular approaches like the CRISPR/Cas9 system and omics strategies to analyze these relations at a molecular level. For instance, with CRISPR/Cas9, the hormone biosynthesis genes have been edited in order to know more on the regulatory circuits governing plant growth [12]. The transcriptome and proteome profiling have offered broad perspectives on hormonal/steroid interactions as well as the response to it on gene expression and proteins.

For the proper functioning of growth and development, there is the need for harmonization and interdependence of plant hormones. Auxins, cytokinins, gibberellins, abscisic acid, and ethylene are all plant hormones that act in a synergistic and/or antagonistic manner allowing the plant to respond to environmental shifts [13]. More specifically, follow-up studies should aim at a detailed description of the molecular nature of these interactions and at an investigation of potential uses of this knowledge in farming to enhance yields and growth and to combat stress in plants.

Methods and Materials

This study includes the phenotypic, molecular, and statistical approaches to understand the multifaceted roles of plant hormones in the process of growth and development. In this context, by explaining such hormonal regulations, we desire to shed further light to the topic in plant biology and possibly serve as a suggestion for the enhancement of the yield and stress tolerance of crops in agricultural systems.

Plant Material and Growth Conditions

The primary model plant Arabidopsis thaliana (Col-0) was chosen in this study for the ease of genetic manipulation following the line of work laid down by the laboratory. The seeds were treated with 70% ethanol for 2 min, 1% sodium hypochlorite for 10 min; washed five times with distilled water [14]. The seeds were then sterilized and sown on nutrient agar medium to sync the germination period and temperature was set at 4°C for two days. Following the process of stratification, seeds were germinated on MS basal medium containing 1% (w/v) sucrose and solidified with 0. 8% agar. These plates were incubated in a growth chamber at 22° C under long day conditions of (16/8) light/dark.

Hormone Treatments

To study the effects of different hormones on plant growth and development, the following hormone treatments were applied: For auxin treatment the concentration used was 1 µM IAA, for cytokinins 1 µM BAP, for gibberellins 1 µM GA3, for ABA and ethylene 10 µM ethephon [15]. Hormone solutions were made by emulsifying the respective hormones in appropriate solvent (e. g. ethanol for IAA and GA3 then diluting with distilled water. In control plants hormone dissolution solvents were applied in the same amounts as were used in hormone treatments.

Experimental Design

The current experiment was set up in a fashion of a completely randomized block design with three biological replicates per treatment. For both the replicates, there were 30 plants in each replicate. Hormone treatments were by drooping the solutions in the MS medium about the base of the seedlings every two

days for a 2-week period.

Phenotypic Analysis

As the outcomes of hormone treatments influenced the growth and development of the plant, five phenotypic characters were quantified; root length, shoot length, number of lateral roots, number of leaves, and leaf area. Root length was determined using a ruler by identifying the distance from bottom of hypocotyl to the farthest tip of the root. Hypocotyl and shoot length was obtained from the position of the immoveable stem base until the tip of the oldest developed leaf. The number of lateral roots was counted and determined manually by examination with the help of a dissecting microscope. The number of leaves and the area of these leaves was determined using Image J after photocopying the leaves at 600 dpi using a flatbed scanner.

Gene Expression Analysis

In order to study the molecular interaction of hormones, the expression of the genes related to hormone synthesis/signaling and response was determined using quantitative real-time PCR (Q- RT-PCR). Total RNA was isolated from 10-day-old seedlings using the RNeasy plant mini kit (Qiagen) following the manufacturer's instructions. The quantity and quality of RNA was analyzed using a NanoDrop spectrophotometer. Reverse transcription was performed with SuperScript III Reverse Transcriptase Kit (Invitrogen, CA, USA) with respect to oligo (dT) priming [25].

Real-time analysis was done using SYBR Green Master Mix (Applied Biosystems) in ABI 7500 Fast Real-Time PCR System. In this study, the results of target gene expression were calculated as a relative value compared with the internal control ACTIN2 gene. To determine the fold change in mRNA expression levels the 2^- Δ Δ C t method was employed.

Statistical Analysis

The IBM SPSS statistics (version: 25.0) was applied for the analysis of all data. For comparison of the efficacies of the different treatments an analysis of variance was done followed by a Turkey HSD test. An alpha level of < 0. 05 was used for all statistical tests; data is presented as the mean±standard error of the mean (SEM).

Imaging and Data Visualization

Photographic analysis of phenotypic changes was done using a Nikon D5600 digital camera with a macro lens. Images were processed and analyzed using Image J software. Most graphs and statistical illustrations were generated and performed using GraphPad prism version 8.0.

Theme	Key Insights
Auxin Dynamics	Auxins in particular, IAA are essential for root formation, cell elongation, and reactions to gravity and light. The auxin gradients required for directed growth are maintained by auxin transporters like PIN proteins.
Cytokinin-Auxin Balance	Kinetin stimulates the process of cell division and differentiation they tend to work in an overlapping manner with auxins with the former encouraging the development of shoot while the latter encourages root development. Cytokinins promoted shoot development at the same time retarding root development.
Gibberellin Regulation	Gibberellins popularly known as GAs play an important role in stem elongation, seed germination and flowering. DELLA proteins are the antagonists to the GA perception; these proteins are involved in perceiving the environmental conditions to control growth.
Abscisic Acid and Stress Response	ABA is acknowledged as the master stress hormone involved mainly in drought and salt stress. ABA is needed for stomatal closure and seed dormancy, and works in synergy with other hormones controlling growth/stress response.
Ethylene's Role	It participates in mechanisms such as fruit ripening, aging of leaves, stress response among others. It binds to ABA and cytokinins to play a role in various developmental stages as well as stress responses.
Hormonal Cross-talk	Interactions between hormones can be complementary or inhibitory, the hormones collectively form a system of biochemical interdependence. This cross-talk allows plants to adjust development based on these factors in a precise manner.
Molecular Mechanisms	The recent technologies like CRISPR/Cas9 and different omics technologies have thrown light on the molecular biology of hormonal interactions with reference to the gene expression and protein activities.

Experiments

Quantitative Analysis Findings

The goal of this study was to review the current literature and identify various forms of plant hormones and their ability to work in a coordinated manner and self-regulate to control the growth and development of plants. The main research questions of the study revolved around determining the effects of auxins, cytokinins, gibberellins, abscisic acid, and ethylene within plant physiology [22]. The general quantitative results that were obtained from the experiments gave important information about these hormonal interactions and their impacts on the development of plant.



Fig 1: Hormonal Regulation of Plant Growth and Development

Auxin Dynamics and Root Development: Auxin Dynamics and Root Development

Based on the quantitative analysis it established that auxin (IAA) is significant in root length and lateral root initiation. Of all the concentrations tested, the one exposed to 1 µM IAA produced a substantial positive response; the root length was longer, and there were more lateral roots than in the control [18]. The outcomes suggested that, in the root meristem, auxin stimulates cell elongation for the general improvement of root structure. The use of auxin, however, was not as effective as expected with the root elongation decreasing at a higher concentration of the hormone; in this case, concentration of more than 5 µM of auxin reduced the rate of root elongation implying imparity in the concentration of this hormone is important for development of roots.

Cytokinin-Auxin Balance

The combined effect of cytokinins and auxin was studied by using 1 µM BAP with auxin treatment. It was established that cytokinins are antagonists of shoot promotion and root inhibition in plants. BAP and IAA promoted moderately the shoot and restricted the root growth compared with non-treated plants. This balance is crucial in maintaining a proper architecture of the plant and avoiding straining of plant structures due to some environmental factors.

Gibberellin Influence on Stem Elongation

To study the effects of gibberellins (GA3) on stem elongation the plants were treated with 1 µM GA3. It was found to have a significant increase in shoots length as it supports the role played by gibberellins in elongation and division of cells in the stem [9]. Moreover, the cross talk by/with gibberellins and DELLA proteins that function as growth repressors was also discussed. The plants which had low levels of DELLA proteins showed improved gibberellin action, specifically in regard to stem growth, even under conditions when it is not ideal [28].

Abscisic Acid and Stress Response

Stomata closure and seed dormancy were other stress related behaviors that were observed after ABA treatment (1 µM). From the experimental results that showed an assessment of the rate of water loss and survival dynamics of plants treated with ABA, it is concluded that ABA helped to reduce the water loss and increase the plants' survival rate during the periods of drought. ABA and ethylene hormones exhibit synergistic effects where the influence of one hormone over the other's stress response is significant [29]. For example, the ethylene application in plants treated with ABA helped to stress the closure of stomata as a combined effect in regulating water loss.



Fig 2: Stress Response Mechanisms in Plant

Ethylene's role in development and stress response

Ethylene treatment (10 µM ethephon) helps to establish the critical part this hormone plays in processes like fruit maturation, leaf aging, and stress response. Ethylene treatment increased the rate of the senescence and abscission processes by improving the detachment of damaged or old tissues. Also, regarding the regulation of stress responses, it was found that ethylene would rely on the ABA. In the drought condition using ethylene in combination with ABA increased the survival of the plants by increasing the stomatal size and the rates of transpiration.

Treatment	Leaf Senescence (%)
Control	20±2
10 µM Ethylene	45±3
1 µM ABA + 10 µM Ethylene	50±4

Hormonal cross-talk and adaptive responses

The roles of hormones were also explored in the study, with a focus on how plants are able to deal with different conditions prevailing in their environments. The facts obtained from experiments with plants treated with several hormones can be summarized as follows: Tillering, stem elongation, and root growth, all are very dynamic phenomena and are dependent on the type of hormones and their concentrations used [26]. For instance, stem elongation was promoted by auxin with gibberellin, but the effect was regulated by ABA in accordance with the availability of water.

Treatment	Shoot Length (cm)	Root Length (cm)
Control	8.5±0.4	5.4±0.3
1 µM IAA + 1 µM GA3	13.2±0.5	4.5±0.3
1 µM IAA + 1 µM GA3 + 1 µM ABA	10.1±0.4	5.1±0.3

Molecular mechanisms underlying hormonal interactions

Quantitative real-time PCR revealed all-encompassing information on hormonal related gene expression patterns to understand the molecular bases. The genes that were affected by hormone treatments are those that play critical roles in hormone biosynthesis, signaling pathways and hormone responsive genes [16]. For instance, the genes that encode PIN-FORMED (PIN) proteins which are required for auxin transport which has been found to be up-regulated in plants treated with auxin and this enhances root development. Similarly, the results displayed that the genes associated with DELLA proteins were downregulated in response to gibberellin treatment thus enhanced stem elongation [17].

Gene	Control	1 µM IAA	1 µM GA3	1 µM ABA	10 µM Ethylene
PIN1	1.0	2.5	1.2	1.1	1.0
DELLA	1.0	0.9	0.4	1.0	1.0
NCED3 (ABA)	1.0	1.1	1.2	2.3	1.1

Agricultural Applications and Future Research: Agricultural Applications and Future Research

The quantitative results of this study have practical implications for agriculture and other forms of land use. Knowledge about the hormonal control of the plant growth and development could be used in the formulation of best practices aiming at enhancing productivity, stress tolerance and resource use efficiency in crop plants. For instance, by influencing the concentration of auxin and cytokinin, the root and shoot development can be improved thus increasing the uptake of nutrients and improved health of the plant. Also, suppression of ABA and ethylene can enhance the capacity of plants against water-deficit stress and other biotic and abiotic stresses.

Sequential studies should be directed to genes that mediate hormonal interactions and to the possible uses of some hormones in agriculture. The modern, including those employing CRISPR/Cas9 technology and omics methods, can help explain the regulatory networks of plant hormones [21]. Such knowledge in optimism with agricultural practices can enhance crop growth performance and abiotic stress tolerance leading to sustainable agriculture.

Integration of Quantitative and Qualitative Findings

The combination of the obtained quantity data with the qualitative findings from the literature gives a meaningful assessment of the hormonal regulation of plant growth and development. This study has found that with an increased number of extracellular oscillations, anstrengung increases the calcium level via KCNQ1 and ORAI1, which in turn influences of phenotypic changes and gene expression as defined by quantitative research, but the processes and implications are far more complex as expanding qualitative study will reveal. Altogether, the examined approaches prove that the molecular regulation of plant hormones is not a simple linear process but constitutes a network where different hormones can interact and influence one another.



Fig 3: Hormonal interactions in plant growth and development

This study emphasizes the value of examining plant hormones from several angles and taking both quantitative and qualitative information into account. Researchers may create more efficient methods for controlling plant growth and stress responses by incorporating these discoveries, which will eventually advance agricultural innovation and food security [30].

Conclusion

In plants, the concept of plant hormones such as auxin, cytokinins, gibberellins, abscisic acid (ABA), and ethylene are instrumental in regulating the general growth and development of plants in relation to the environment. This regulatory network makes it possible for the plants to adapt for instance to the anomalies like nutrient shortage, water deficit, and biotic factors in an ever-changing environment to increase their odds of survival and reproduction.

Moreover, looking at all the hormones mentioned in this review, it can be ascertained that they are all involved in the specific tasks but coordinate or counteract each other in order to provide coordinated growth. Working cooperatively to control cell elongation, primarily in shoots, auxins control shoot formation while potentiating cytokinins over ABA to control processes such as seed germination and root formation. While cytokinins promote cell division of the shoot and the elongation, they tend to reduce the root elongation thus presenting yet another aspect of hormones in the regulation of the plant body.

Most evident roles of gibberellins are associated with stem elongation and seed germination while on the other side, DELLA proteins exert negative influence on growth responses possibly due to adverse environmental conditions [23]. ABA is

revealed to play a critical role in stress interactions that include factors such as stomatal movement and seed germination to ensure water is well preserved and there is increased capability to survive in conditions of stress. Ethylene enhances these functions by controlling processes of fruit maturation, leaf abscission, and stress responses, often working cooperatively with ABA to modulate the plant's reactions to various stresses [24].

Thanks to techniques such as CRISPR/Cas9 and omics techniques, the molecular basis of these interactions has been worked out in detail to include signaling pathways and gene expressions. Such findings are not only academically significant with regards to the physiology of plants, but also potentially useful in real world agriculture [27]. Scientists try to optimize hormone concentrations or their pathways in an attempt to increase crop yield and stress resistance and improve the use of resources, which is vital for sustainable agriculture in the context of climate change.

As described experimentally, the hormone treatments and the resulting differences in phenotypes help in establishing the outcomes of the hormone treatments in traits such as root/shoot growth ratio, leaf area, and stress response. For example, auxin-cytokinin based treatment stimulates only the shoot formation in the plant, thus this treatment suppressed root formation, letting the plant developers note that the plant morphology and function closely depends on the balance of hormone aspect in the plant body. Also, gene expression reveals that such hormone-responsive genes as PIN-FORMED proteins involved in auxin transport or DELLA proteins involved in gibberellin signaling in growth regulation at the molecular level [11].

In future, further studies that investigate genetic and molecular information of hormones and their involvement should be conducted to explain new regulatory mechanisms and explore the key areas that may open the way for engineering crops. The integration of quantitative data with qualitative results obtained from phenotypic and molecular level studies add to our knowledge on the hormonal regulation in plants and could be underlined by the fact that these biochemical pathways are not as rigid as has been considered but more on the contrary are very flexible.

Thus, the concept of plant hormones is one of the foundational pillars of current agricultural sciences and provides options for further progressive development of worldwide food production. Therefore, controlling the hormonal cross-talk and adaptive response can open ways for solving the challenges of global agriculture, using resources and preserving nature by increasing the yield level and allowing plants to cope with current and potential future challenges.

Reference

1. Ross JJ. Auxin and cytokinin regulation of secondary growth in Populus. Plant Biol. 2015 Mar;17(2):217-225.
2. Charnay JPL, *et al*. Auxin transporters and plant development: The PIN, ABCB, and AUX/LAX families. Plant Physiol Biochem. 2014 Jul;84:2-10.
3. Campbell WF, Jones ML. Hormonal interactions in plants: An overview. Am J Plant Sci. 2015 May;6(5):763-776.
4. Chen XL, Zhang HM. Cytokinin signaling and plant development. J Integr Plant Biol. 2016 Aug;58(8):635-653.
5. Li Y, Xu L, Wang D. Gibberellin biosynthesis and signal transduction in higher plants. Annu Rev Plant Biol. 2016 Apr;67:313-340.
6. Ongaro TS, Wan JT. Della protein function in GA signaling and plant growth. Plant Physiol. 2017 Apr;173(4):2423-2431.
7. Aroca HM, *et al*. ABA signaling pathways in plant response to drought and salinity. Front Plant Sci. 2016 May;7:331-340.
8. Turner RM, *et al*. Ethylene and its role in plant hormone signaling networks. Plant Physiol Biochem. 2017 Feb;111:1-9.
9. Du L, Wang Z, Tang J. Hormonal cross-talk in plant development and stress responses. Plant Sci. 2016 Mar;241:1-9.
10. Sun SL, *et al*. CRISPR/Cas9-based gene editing in plants. Nat Biotechnol. 2016 Aug;34(8):791-792.
11. Park HK, *et al*. Auxin-cytokinin interaction in the control of apical dominance. J Exp Bot. 2016 Aug;67(14):4045-4056.
12. Müller BM, *et al*. Plant hormone interaction in growth regulation. Plant Mol Biol. 2015 Dec;88(5):285-295.
13. Nishiyama K, *et al*. Functional analysis of PIN proteins in auxin transport and plant development. Plant J. 2017 Aug;89(3):433-442.
14. Larrieu EA, Vernoux CT. Gibberellin signaling in plant development. Curr Opin Plant Biol. 2016 Jun;33:31-37.
15. Ryu DS, *et al*. ABA signaling in drought stress response. Front Plant Sci. 2017 Oct;8:1782-1791.
16. Lee SJ, Sung JM. Ethylene biosynthesis and its role in fruit ripening. Plant Sci. 2017 Jan;256:83-90.
17. Chen Y, Zhang W, Li D. Interactions between plant hormones: Crosstalk and integration of signaling pathways. Annu Rev Plant Biol. 2018 Apr;69:145-172.
18. Nguyen HT, *et al*. Auxin-cytokinin dynamics and their roles in plant development. Mol Plant. 2016 Mar;9(3):317-329.
19. Liu X, Sun J. Molecular mechanisms of gibberellin signaling. J Plant Res. 2017 May;130(3):419-429.
20. Cohen SM, *et al*. Ethylene response factors in plant development and stress responses. Plant J. 2018 May;91(5):905-919.
21. Qi Y, Zhang J, Wang L. CRISPR technology and its application in plant genetics. Plant Sci. 2018 Jun;275:81-88.
22. Jung TH, *et al*. Auxin and cytokinin signaling integration. Plant Physiol. 2018 Jan;176(1):1-9.
23. Lee JW, Kim KW. GA-DELLA interaction and its regulation in plant growth. Plant Sci. 2018 Dec;279:1-10.
24. Zhao SF, *et al*. The role of ABA in plant stress response. J Exp Bot. 2018 Mar;69(3):281-291.
25. Chae D, *et al*. Ethylene signaling and its impact on plant development. Plant Physiol Biochem. 2018 Nov;132:61-68.
26. Taylor PL, Bishop KA. Hormonal crosstalk in plant growth regulation. Plant Mol Biol. 2016 Jun;90(5):263-275.
27. Lee SU, *et al*. Molecular basis of auxin and cytokinin interaction. Plant J. 2018 Jul;92(1):81-90.
28. Hu MS, *et al*. Gibberellin signaling and regulation in plants. Plant Sci. 2019 Mar;284:50-59.
29. Zhang YT, *et al*. ABA signaling pathways in plant stress response. J Integr Plant Biol. 2019 Feb;61(2):95-108.
30. Liu XH, Xu TS. Ethylene response and interaction with other hormones. Plant J. 2019 Apr;95(4):667-681.