



ISSN (E): 2320-3862
ISSN (P): 2394-0530
NAAS Rating: 3.53
JMPS 2018; 6(1): 27-30
© 2018 JMPS
Received: 07-11-2017
Accepted: 08-12-2017

Swati Celia Topno
P.G. Dept. of Botany,
Kolhan University, Chaibasa,
West Singhbhum, Jharkhand,
India

Manoj Ranjan Sinha
HOD, Department of Botany,
Co-operative College,
Jamshedpur, Jharkhand, India

Phytochemical analysis of floral extract of three medicinal plants used by local inhabitants of West Singhbhum district, Jharkhand (India) to cure skin diseases

Swati Celia Topno and Manoj Ranjan Sinha

Abstract

In the present study floral extract of three medicinal plants viz., *Acacia nilotica* (Linn.) Delile, *Butea monosperma* (Lam.) Taub. And *Woodfordia fruticosa* Kurz. with aqueous, ethanol and methanol extract were screened for qualitative phytochemical analysis. Qualitative phytochemical analysis was performed for the test of proteins, carbohydrates, phenol-tannins, flavonoides, saponin, glycosides, steroids, phlobatannins, alkaloids and terpenoides with the help of standard protocol.

The qualitative phytochemical analysis revealed the presence of protein, carbohydrates, phenol and tannin, saponin, glycosides, steroids, alkaloids and terpenoids in extracts of flower of The results suggest that flowers of all the above mentioned plants. On the basis of presence of active phytochemical, the present paper provide justification for the use of medicinal plants by the local inhabitants of West Singhbhum district, Jharkhand (India) to cure skin diseases.

Keywords: Biological diversity, tribal, medicinal plants, skindiseases, phytochemical analysis

Introduction

Medicinal plants have been used to treat human diseases for thousands of years because they have vast and diverse assortment of organic compounds that can produce a definite physiological action on the human body. The skin is a vital organ of the human body, which is characterized by the largest possible surface area, and it has a direct contact with the environment. Skin diseases in developing countries have a serious impact on people's quality of life and bring out significant burden to the nations. It affects more than 60% of the general population [1]. Skin disease refers to disorders of predominantly the superficial layers of the skin. The prevalence of skin disease in any region or country depends on various factors, such as genetics, racial constitution, social and hygienic standards, customs and occupations. Transmissible skin diseases are observed in people who are living under poor socioeconomic and unhygienic conditions [2]. In India there is a significant incidence of infectious disorders in rural communities because of underdeveloped economy and social backwardness [3]. Ignorance on seriousness of the disease and improper medication worsens the condition in the rural areas of West Singhbhum where most of the tribal are engaged in agriculture and mining. Medicines can induce skin reactions and some are potentially life threatening [4]. Up to 80% of the population suffering from skin problems may not seek medical help [5]. The common skin problems are Acne, Burn, scars, Psoriasis, Scabies, Skin grafting, Vitiligo, Pediculosis, Herpes simplex infection, Varicella, Herpes Zoster, Erythema, Urticaria etc. They are found in children, young and adults as well as in old persons. Usually for peak level skin disorder, the therapy of skin problems is longer for complete removal of problems. In all over the world use of drug like Benzoyl Peroxides, Proactive, Antibiotics, Retin-A, Oral retinoid, Salicylic acid, Anti-Histaminic, Minerals and Vitamins, Steroids, Analgesic are of more interest for skin specialist for the modern treatment. But the herbal medicine is becoming popular due to toxicity and side effects of allopathic medicines [6].

Most important of such compounds are alkaloids, tannins, flavonoids, terpenoids, saponins and phenolic compounds. Pharmacists are interested in these compounds because of their therapeutic performance and low toxicity [7].

Correspondence

Swati Celia Topno
P.G. Dept. of Botany,
Kolhan University, Chaibasa,
West Singhbhum, Jharkhand,
India

Due to their natural origin and low toxicity, phenolic compounds are a promising tool in eliminating the causes and effects of skin aging, skin diseases, and skin damage, including wounds and burns. Hence, present paper emphasize on evaluation and characterization of three plants viz. *Acacia nilotica* (Linn.) Delile and *Butea monosperma* (Lam.) Taub. belonging to Fabaceae family and *Wood fruticosa* Kurz. belongs to Lythraceae family [8] and plant constituents against a number of skin diseases based on their traditional claims of the plants. (figure 1, 2 and 3 respectively).



Fig 1: *Acacia nilotica* (Linn.) Delile figure



Fig 2: *Butea monosperma* (Lam.) Taub



Fig 3: *Woodfordia fruticosa* Kurz

Materials and Methods

The plant materials were shade dried until all the water molecules evaporated and plants became well dried for grinding. After drying, the plant materials were ground well using mechanical blender into fine powder and transferred into airtight containers with proper labeling for future use.

Preparation of plant extracts

Hot water extraction: 5gm of dried finely powdered plant material was taken in a beaker and 200ml of distilled water was added. The mixture was heated on a hot plate with continuous stirring at 30°-40°C for 20 minutes. Then the water extract was filtered through filter paper and the filtrate was used for the phytochemical analysis. The water extract was kept in refrigerator when not in use.

Solvent extraction

Crude plant extract was prepared by Soxhlet extraction method. About 20gm of powdered plant material was uniformly packed into a thimble and extracted with 250ml of different solvents separately. Solvents used were methanol and ethanol. The process of extraction continues for 24 hours or till the solvent in siphon tube of an extractor become colorless. After that the extract was taken in a beaker and kept on hot plate and heated at 30-40°C till all the solvent got evaporated. Dried extract was kept in refrigerator at 4°C for their future use in qualitative phytochemical analysis. The extract was tested for the presence of bioactive compounds by using following standard methods [9, 10, 11].

Test for proteins

Millon's test

Crude extract when mixed with 2ml of Millon's reagent, white precipitate appeared which turned red upon gentle heating that confirmed the presence of protein.

Ninhydrin test

Crude extract when boiled with 2ml of 0.2% solution of Ninhydrin, violet colour appeared suggesting the presence of amino acids and proteins.

Test for carbohydrates

Fehling's test

Equal volume of Fehling A and Fehling B reagents were mixed together and 2ml of it was added to crude extract and gently boiled. A brick red precipitate appeared at the bottom of the test tube indicated the presence of reducing sugars.

Benedict's test

Crude extract when mixed with 2ml of Benedict's reagent and boiled, a reddish brown precipitate formed which indicated the presence of the carbohydrates.

Iodine test

Crude extract was mixed with 2ml of iodine solution. A dark blue or purple coloration indicated the presence of the carbohydrate.

Test for phenols and tannins

Crude extract was mixed with 2ml of 2% solution of FeCl₃. A blue-green or black coloration indicated the presence of phenols and tannins.

Test for flavonoids

Shinoda test

Crude extract was mixed with few fragments of magnesium ribbon and concentrated HCl was added drop wise. Pink scarlet colour appeared after few minutes which indicated the presence of flavonoids.

Alkaline reagent test

Crude extract was mixed with 2ml of 2% solution of NaOH. An intense yellow colour was formed which turned colourless

on addition of few drops of diluted acid which indicated the presence of flavonoids.

Test for saponins

Crude extract was mixed with 5ml of distilled water in a test tube and it was shaken vigorously. The formation of stable foam was taken as an indication for the presence of saponins.

Test for glycosides

Liebermann's test

Crude extract was mixed with each of 2ml of chloroform and 2ml of acetic acid. The mixture was cooled in ice. Carefully concentrated H₂SO₄ was added. A colour change from violet to blue to green indicated the presence of steroidal nucleus, i.e., glycone portion of glycoside.

Salkowski's test

Crude extract was mixed with 2ml of chloroform. Then 2ml of concentrated H₂SO₄ was added carefully and shaken gently. A reddish brown colour indicated the presence of steroidal ring, i.e., glycone portion of the glycoside.

Keller-kilani test

Crude extract was mixed with 2ml of glacial acetic acid containing 1-2 drops of 2% solution of FeCl₃. The mixture was then poured into another test tube containing 2ml of concentrated H₂SO₄. A brown ring at the interphase indicated the presence of cardiac glycosides.

Test for steroid

Liebermann's Test

2 ml Crude extract was mixed with 2 ml acetic acid (CH₃COOH) and then 1 ml of concentrated H₂SO₄ was added

drop wise, the presence of blue green color indicates the presence of steroids.

Salkowski Test

2 ml Crude extract was mixed with 2ml of chloroform and concentrated H₂SO₄ was added sidewise. A red colour produced in the lower chloroform layer indicated the presence of steroids.

Liebermann's Bruchard Test

2 ml Crude extract is mixed with 2ml of chloroform. Then 2ml of each of concentrated H₂SO₄ and acetic acid were poured into the mixture. The development of a greenish coloration indicated the presence of steroids.

Test for Phlobatannin

2 ml Crude extract is mixed with 2ml of 1% HCl which gives red precipitate on gentle heating confirms the presence of phlobatannin.

Test for alkaloids

2 ml Crude extract was mixed with 2ml of 1% HCl and heated gently. Mayer's And Wagner's reagents were then added to the mixture. Turbidity of the resulting precipitate was taken as evidence for the presence of alkaloids.

Test for terpenoids

Crude extract was dissolved in 2ml of chloroform and evaporated to dryness. To this, 2ml of concentrated H₂SO₄ was added and heated for about 2 minutes. A grayish colour indicated the presence of terpenoids.

Result and Discussion

Table 1: Qualitative phytochemical analysis of floral extract of three medicinal plants

S. No	Phytochemical test	<i>Acacia nilotica</i> (Linn.) Delile			<i>Butea monosperma</i> (Lam.) Taub.			<i>Woodfordia fruticosa</i> Kurz.		
		A.E	E.E	M.E	A.E	E.E	M.E	A.E	E.E	M.E
1	Proteins	+	+	+	+	+	+	-	+	+
2	Carbohydrates	+	+	+	+	+	+	+	+	+
3	Phenol & Tannins	+	+	+	+	+	+	+	+	+
4	Flavonoids	+	-	+	-	-	+	+	+	+
5	Saponin	-	-	+	+	+	+	+	+	+
6	Glycosides	+	+	+	+	+	+	+	+	+
7	Steroids	+	+	+	+	+	+	-	+	+
8	Phlobatannins	-	-	+	-	-	-	-	-	-
9	Alkaloids	-	+	+	+	+	+	-	+	+
10	Terpenoids	-	-	-	-	+	+	+	+	+

A.E= Aqueous extract, E.E = Ethanolic extract, M.E= Methanolic extract, (+) = Present, (-) = Absent

Result

- In *Acacia nilotica* (Linn.) Delile it has been observed that proteins, carbohydrate, phenol & tannins, glycosides and steroids were present in all the three extracts whereas saponin and phlobatannins were found only in methanolic extract. Flavonoids are present only in ethanolic plant extract. Alkaloid was found absent in aqueous extract. It was observed that terpenoids were absent in all three plant extract.
- In *Butea monosperma* (Lam.) Taub. Proteins, carbohydrates, phenol & tannins, saponin glycosides, steroids and alkaloids were observed to be present in all the three extract while it was found that flavonoid was only present in methanolic extract. Phlobatannin was absent in all the three extracts and it was observed that terpenoids was present in ethanolic and methanolic plant extracts.

- In *Woodfordia fruticosa* Kurz. Carbohydrate, phenol & tannins, flavonoids, saponins were observed to be present in all the three extracts whereas protein, steroids and alkaloids were present in aqueous extract and it was found that phlobatannin was absent in all the three extracts.

Discussion

From the result obtained by the phytochemical analysis of three medicinal plants, it could be seen that carbohydrates, phenol & tannins and glycosides were present in all the three plants. Several reports revealed that medicinal plants are rich in phenolic compounds and have antioxidant properties [12, 13]. (Phenolic compounds also possess potent antifungal, antiviral and antibacterial activity [14]. It is also mentioned that phytochemical analysis on plants extracts revealed the presence of constituents that are known to exhibit medicinal as well as physiological activities [9]. The plant extracts also

revealed to contain phenol & tannins. It was reported that tannins contribute property of astringency i.e. fasten the healing of wounds and inflamed mucous membrane and have received considerable attention in the fields of nutrition, health and medicine, largely due to their physiological activity, such as antioxidant, antimicrobial and anti-inflammatory properties^[15]. Flavonoids were found present in methanolic plant extract of all the three plants. Several studies shows that flavonoids are a group of polyphenolic compounds which influence the radical scavenging, inhibition of hydrolytic and oxidative enzymes and also act as anti-inflammatory agent^[16]. Saponins was found present in methanolic plant extract of *Acacia nilotica* (Linn.) Delile while it was found in all the three plant extract of *Butea monosperma* (Lam.) Taub. and *Woodfordia fruticosa* Kurz. According to several studies the plant extracts were also revealed to contain saponins which are known to produce inhibitory effect on inflammation^[17]. Steroids have been reported to have antibacterial properties^[18]. Terpenoids are active against bacteria^[19, 20, 21].

Acknowledgement

The first author is highly grateful to Dr. KiranShukla, Head, P.G. Department of Botany, Kolhan University, Dr. Vishnu Shankar Sinha, Head, Department of Botany, Tata College Chaibasa for providing their valuable suggestions and Mrs K Boipai, Principal, Tata College, Chaibasa for providing college science laboratory to carry on the phytochemical analysis for my research work. The first author is also grateful to UGC for financial assistance in the form of fellowship under the scheme of NFHE (National Fellowship for Higher Education for ST candidates).

References

- Kocinaj A, Kocinaj D, Berisha M.; Skin disease among preschool children. J Bacteriol. Res. 2009; 1(2):25-29.
- Al-saeed WY, Al-Dawood Bukhari KM, Bahnassy IA. Risk factors and co morbidity of skin disorders among female school children in eastern Saudi Arabia. Invest Clin. 2007; 48(2):199-212.
- Dayal SG, Gupta GD. A cross section of skin diseases in Bundelkhand region, UP. Ind. J Derm. Ven. Leprol. 1972; 43:258-26.
- Lee A, Thompson J. Drug induced skin reactions. Adverse Drug Reactions, 2nd edition, Pharmaceutical Press. 2006, 125-156.
- Williams HC. Epidemiology of skin diseases. Rook's Textbook of Dermatology. 7th edition, Oxford: Blackwell Science. 2004; 6.1-6.21.
- Samraj K, Thillaivanan S, Parthiban P, Samraj K. A review of beneficial effects of medicinal plants on skin and skin diseases. 2014; 3(1):93-106.
- Inayatullah S, Prenzler PD, Obied HK, Rehman AU, Mirza B. Bioprospect-ing traditional Pakistani medicinal plants for potent antioxidants Food. 2012; 132:222-229.
- Khare CP. Indian Medicinal Plants: An Illustrated Dictionary. 2007; 4:105.719.
- Sofawora A. Medicinal Plants And traditional Medicine in Africa. Spectrum Books Ltd., Ibadan, Nigeria. 1993,191-289.
- Trease GE, Evans WC. Pharmacognosy, 11th edn., Bailliere Tindall, London. 1989, 45-50.
- Harborne JB. Phytochemicals Methods. Chapman and Hall Ltd, London. 1973, 49-188.
- Brown JE, Rice-Evans CA. Luteolin rich artichoke extract protects low density Lipoprotein from oxidation in vitro. Free Radical Res. 1998; 29:247-255.
- Krings U, Berger RG. Antioxidant activity of roasted foods. Food Chem. 2001; 72:223-229.
- Czemplik M, Zuk M, Kulma AS, Szopa J. GM flax as a source of effective Antimicrobial compounds. Sci. Microb. Pathog. Commun. Curr. Res. Technol. Adv. 2011; 76:39-47.
- Santos-Bulega C. Scalbert AJ. Proanthocyanidins and tannin – like Compounds – Nature, occurrence, dietary intake and effects on nutrition and health. Sci Food Agric. 2000; 80:1094-1117.
- Frankel E. Nutritional benefits of flavonoids, International conference on food factors: chemistry and cancer prevention, Hamamatsu, Japan, Abstract, C-2, 1995.
- Just MJ, Recio MC, Giner RM, Cueller MU, Manez S, Billia AR, Rios JL. Antimicrobial activity of unusual lupine saponins from *Bupleurum fruticosens*. 1998; 64:404-407.
- Raquel FE. Bacterial lipid composition and antimicrobial efficacy of cationic steroids compounds. Biochemicaet Biophysica Acta. 2007; 2500-2509
- Barre JT, Bowden FB, Coll JC, Jesus J, Fuente VE, Janairo GC. A bioactive triterpene from *Lantana camara*. Phytochemistry. 1997; 45:321-324.
- Habtemariam S, Gray AI, Waterman PG. A new antibacterial sesquiterpene from *Prema oligotricha*. Journal of Natural products. 1993; 56:140-143.
- Scortichini M, Pia RM. Preliminary in vitro evaluation of the antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora* (Burkill) Winslow. Journal of Applied Bacteriology. 1991; 71:109-112.
- Frankel E. Nutritional benefits of flavonoids, International conference on food factors: chemistry and cancer prevention, Hamamatsu, Japan, Abstract, C-2 15.
- Santos- Bulega C., Scalbert, A.J. 2000, Proanthocyanidins and tannin – like Compounds – Nature, occurrence, dietary intake and effects on nutrition and health. Sci Food Agric. 1995; 80:1094-1117.
- Just MJ, Recio MC, Giner RM, Cueller MU, Manez S, Billia AR, *et al.* Antimicrobial activity of unusual lupine saponins from *Bupleurum fruticosens*. 1998; 64:404-407.
- Barre JT, Bowden FB, Coll JC, Jesus J, Fuente VE, Janairo GC. A bioactive triterpene from *Lantana camara*. Phytochemistry. 1997; 45:321-324.
- Habtemariam S, Gray AI, Waterman PG. A new antibacterial sesquiterpene from *Prema oligotricha*. Journal of Natural products. 1993; 56:140-143.
- Scortichini M, Pia RM. Preliminary in vitro evaluation of the antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora* (Burkill) Winslow. Journal of Applied Bacteriology. 1991; 71:109-112.
- Raquel FE. Bacterial lipid composition and antimicrobial efficacy of cationic steroids compounds. Biochemicaet Biophysica Acta. 2007, 2500-2509.