



ISSN (E): 2320-3362
ISSN (P): 2394-0530
www.plantsjournal.com
JMPS 2021; 9(5): 24-27
© 2021 JMPS
Received: 16-06-2021
Accepted: 24-08-2021

M Saminathan

PG & Research,
Department of Botany, Mannai
Rajagopalaswamy Government
Arts College, Mannargudi 614
001, Tamil Nadu affiliated to
Bharathidasan University,
Tiruchirappalli, India

Dr. AC Tangavelou

Department of Plant Biology
and Plant Biotechnology, Guru
Nanak College (Autonomous),
Chennai, Tamil Nadu, India

A Murugananadham

PG & Research Department of
Botany, Mannai
Rajagopalaswamy Government
Arts College, Mannargudi 614
001, Tamil Nadu affiliated to
Bharathidasan University,
Tiruchirappalli, India

Corresponding Author:

Dr. AC Tangavelou

Department of Plant Biology
and Plant Biotechnology, Guru
Nanak College (Autonomous),
Chennai, Tamil Nadu, India

Ethnomedicinal plants used by the Malayali Tribals of Pachamalai hills for wounds

M Saminathan, Dr. AC Tangavelou and A Murugananadham

Abstract

In the present study, ethnomedicinal surveys were carried out among the Malayali tribes of Pachamalai hills in Tamil Nadu and documented the traditional knowledge on medicinal plants used by them for human health care with special reference to wounds. Totally 26 plant species belonged to 18 families and 26 genera were used for treating wounds by Malayali tribes residing in Pachamalai hills. All the documented information on ethnomedicinal plants for wounds were given alphabetically with correct Botanical name, family, local name, herbarium specimen, habit, flowering & fruiting, uses, dosage of preparation, method of administration along with the recommended diet prescribed by them. This study would provide a firsthand resource material and very useful for the ethnopharmacologist to develop herbal based antidote drug(s).

Keywords: Ethnomedicine, Malayali tribes, Pachamalai hills, Wounds, Drug Discovery

1. Introduction

Ethnomedicine is concerned with the cultural interpretations of health, disease and illness and also addresses the health care-seeking process and healing practices (Pieroni *et al.*, 2005) [19]. According to the World Health Organization, Ethnomedicine maintained its popularity in all regions of the developing world and its use is rapidly expanding in the industrialized countries (WHO, 2003) [21]. Ethnomedicine, a study of traditional medicine, is significant in drug discovery and development for the prevention, treatment and management of novel pandemic disease Covid-19 (Adeleya *et al.*, 2021) [1]. It is stated that around 25% of the modern remedies used globally have been obtained from medicinal plants (Gemechu *et al.*, 2021) [8]. Western trained physicians could not ignore the impact of ethnomedicine on their patients and the research interest in the area of ethnomedicine have been increased tremendously in the last decade (Williams, 2006) [22]. Since the inception of the discipline, scientific research in ethnomedicine has made important contribution to the understanding of traditional subsistence, medical knowledge and practice.

It is interesting to note that the ethnomedicinal uses of plants is one of the most successful criteria used by the pharmaceutical industry in finding new therapeutic agents for the various fields of biomedicine (Cox and Balick, 1994) [4]. Some outstanding medicinal drugs which have been developed from the ethnomedicinal uses of plants include: vinblastine and vincristine from *Catharanthus roseus* (the periwinkle) used for treating acute lymphoma, acute leukaemias etc., reserpine from *Rauwolfia serpentina* (Indian snake root) used for treating hypertension, aspirin from *Salix purpurea* (willow) used for treating inflammation, pain and thrombosis and quinine from *Cinchona pubescens* (cinchona) used for treating malaria. Based on the ethnomedicine, a total of 122 compounds were identified and 80% of these compounds were used for the same (or related) ethnomedicinal purposes (Fabricant and Farnsworth, 2001) [5]. Further, it was discovered that these compounds were derived from only 94 species of plants (Farnsworth and Soejarto, 1985) [6]. Hence it is very urgent to document the knowledge on several other plants from indigenous people, the treasurer of plant wealth. With this background the present study deals with the documentation of ethnomedicinal information on plants from the Malayali tribals of Pachamalai hills and presented here with special reference to wounds.

2. Materials and methods

2.1 Study Area

Pachaimalai Hills, also known Green hill (*Pachai* means green in Tamil), are Low Mountain

range in the Eastern Ghats located between the Salem and Tiruchirappalli districts of Tamil Nadu, South India. It lies between 78.31' East longitude and 11.28' North latitude at an altitude of 2000 to 3000 feet above mean sea level. Temperature is moderately high and the average temperature during summer is 34 °C and lesser in winter. Average humidity ranges from 31 to 33% during November to December. Annual rainfall ranges from 850 to 910 mm. Soil is shallow in rocky areas and deeper in valleys with little or no humus. Vegetation is Scrub jungles mixed with tropical dry evergreen forest and tropical dry deciduous type.

2.2 Malayali tribals

Malayali tribals residing at Achinadu (Gonganavalli), Kombai, Perumal Kovil, Thenpuranadu, Vannadu areas of Pachamalai regions performing agricultural practices and also collect tuber, honey and perform fishing operations in the reserve for their sustenance. At present they do not hunt animals.

2.3 Ethnomedicinal Survey

Intensive field ethnomedicinal surveys were conducted in the Pachamalai hills from 2018 to 2020 to document ethnomedicinal plants resources used by Malayali tribals residing in Pachamalai hills settlements such as Achinadu (Gonganavalli), Kombai, Perumal Kovil, Thenpuranadu, Vannadu regions by following the methods of Jain (1981) [14]. The questionnaires were used to obtain information on medicinal plants with their local names, parts used as medicine, mode of preparation and dosage of administration. The species mentioned by the tribes were taxonomically identified. While gathering the information, herbarium specimens were prepared for all the plant species on ethnomedicinal plants available in the study area by following the methods of Jain and Rao (1976) [13] and deposited at the Department of Botany, Mannai Rajagopalaswamy Government Arts College, Mannargudi, for reference. The herbarium specimens were botanically confirmed by using various regional Flora such as Gamble and Fischer (1915-1936) [7], Matthew (1981-1988) [17] while the Flora of Tamil Nadu by Henry *et al.*, (1987 & 1989) [10, 9] and Nair and Henry (1983) [18] were used to refer the correct botanical names.

3. Results and discussion

In the present study, totally 26 species belonging to 18 families and 26 genera of ethnomedicinal plants were used by the Malayali tribals of Pachamalai hills specifically for the treatment of wounds. Among habitwise, herbs are the dominant habit form represented by 14 species followed by trees (6 species), shrubs (3 species) and climbing herb, liana and small tree (single species each) respectively.

All the documented information on ethnomedicinal plants for the treatment of wounds used by Malayali tribes from Pachamalai hills were given below alphabetically with correct Botanical name, family, local name, habit, uses, dosage and method of treatment.

***Abrus precatorius* L.**, Fabaceae, *Kundumani*, Climbing herb

- Leaf is made into paste, mixed with coconut oil is applied externally for skin diseases and fresh wounds.

***Acacia leucophloea* (Roxb.) Willd.**, Mimosaceae, 'Velaamaram', Tree.

- Leaf is ground into fine paste is applied externally for cuts and wounds for 3 days.

***Acalypha indica* L.** Euphorbiaceae, *Kuppaimeni*, Herb

- Leaf is ground into paste, mixed with coconut oil is applied externally for cuts /wounds.

***Achyranthes aspera* L.**, Amaranthaceae, *Naayurivi*, Herb

- Leaf paste is made into paste along with coconut oil is applied externally for skin itching, wound healing.
- Leaves crushed by hand to extract the juice and the juice applied for wounds.

***Aerva lanata* (L.) Juss. ex Schultes**, Amaranthaceae, *Poolaapoo*, Herb

- Fresh leaf paste mixed with coconut oil is applied externally on wounds.

***Boerhavia diffusa* L.**, Nyctaginaceae, *Saranaichedi*, Herb

- Leaves made into a paste along with coconut oil is applied externally for healing wounds.

***Clausena dentata* (Willd.) M. Roem.** Rutaceae, *Kaattukaruveppilai*, Shrub

- Leaf is made into paste is applied externally on wounds for healing quickly.

***Commelina benghalensis* L.**, Commelinaceae, *Kaanaavazhai /Mayilkaanaa*, Herb

- Whole plant is boiled with water and filtered the decoction is taken three times a day for 4-5 days to cure ear wounds.

***Coscinium fenestratum* (Gaertn.) Coleb.**, Menispermaceae, *Maramanjai*, Liana

- Wood ground into a paste is mixed with coconut oil and is applied on the forehead for quick relief of headache, migraine and also to heal cuts and wounds.

***Datura innoxia* Mill.** Solanaceae, *Oomaththai*, Shrub

- Leaf paste mixed with coconut oil is applied for cuts, wounds, boils and burns.

***Emilia sonchifolia* (L.) DC.**, Asteraceae, *Arukuruvi pacchilai, Muyalchevi*, Herb

- Fresh leaf paste is applied externally for healing fresh cuts and wounds.

***Eupatorium odoratum* L.** Asteraceae, *Neththichedi*, Herb

- Fresh Leaf paste is applied externally directly on the wounds for quick healing.

***Ficus racemosa* L.**, Moraceae, *Aththi*, Tree

- White milky latex is applied directly on fresh wounds for quick healing.

***Leucas aspera* (Willd.) Link**, Lamiaceae, *Thumbai*, Herb

- Leaf made into a paste and mixed with coconut oil is applied externally on wounds.

***Mallotus philippensis* (Lam.) Muell. Arg.**, Euphorbiaceae, *Sokkalaamaram*, Tree

- Leaf made into a paste and mixed with coconut oil is applied externally to stop bleeding from wounds.

***Moringa oleifera* Lam.**, Moringaceae, *Murungamaram*, Tree.

- Stem bark and young root ground into a paste along with 5 g salt and apply at throat to cure internal wounds.

Phyllanthus amarus Schumach & Thonn, Euphorbiaceae, *Keezhanelli*, Herb.

- Leaves boiled with water and the filtered decoction is taken thrice a day for the treatment of fresh wounds or any swollen wounds.

Ruellia tuberosa L., Acanthaceae, '*Vedimuthu/ Vettupoondi*', Herb.

- Leaves mixed with 2 small onion, ground into a fine paste and applied on the chronic wounds to cure.

Sida acuta Burm.f., Malvaceae, *Chinnakurunthoati*, Herb

- Leaves boiled with water and the filtered decoction is taken thrice a day for the treatment of breaking of wounds or any swollen wounds.

Solanum torvum Sw., Solanaceae, '*Sundakachedi /Sundaangkodi*', Shrub.

- Leaf made into paste mixed with coconut oil and paste by adding little water and apply it on cuts and wounds to cure.

Sphaeranthus indicus L. Asteraceae, *Kottaikarandhai*, Herb

- Leaf made into paste from one handful of leaves mixed with coconut oil is applied externally for the treatment of skin itching, wounds.

Tamarindus indica L., Caesalpiniaceae, *Puliyamaram*, Tree.

- Stem bark made into a powder and mixed with coconut oil and applied on the wounds for 4 days, till it cures.

Tridax procumbens L. Asteraceae, *Vettukayathazhai*, Herb

- Fresh leaf juice is applied externally as a tincture for fresh burns, cuts and wounds.

Withania somnifera (L.) Dunal, Solanaceae, *Amukkurachedi*, Herb

- Root is made into paste by adding a little amount of water and applied it on wounds for 2 times a day to cure.

Wrightia tinctoria R.Br. Apocynaceae, *Vepaalaimaram*, Tree

- Leaves along with 1 piece of onion and 2 pepper, is made into and applied externally to cure the wounds.

Ximenia Americana L., Olacaceae, *Sengaathirimaram*, Small tree

- Stem bark powder is made into paste by mixing with goat's milk and the paste is applied externally on burn or fresh wounds.

Tribals in the adjacent hills also used these plants for wounds such as *Acacia leucophloea* and *Clausena dentata* used for wounds by the Paliyan tribals of Madurai district (Ignacimuthu and Ayyanar, 2006). These species contain valuable chemical substances and are useful to cure wounds. Wound healing involves a complex interaction between epidermal and dermal cells, the extra cellular matrix, controlled angiogenesis and plasma-derived proteins all coordinated by an array of cytokines and growth factors (Bhat *et al.*, 2007) [3]. To provide evidence for this present study, most of the plants are used scientifically proved for wound healing activity such as *Tridax procumbens* (Udupa *et al.*, 1991; Babu *et al.*, 2003; Johnson and Gorle, 2012) [20, 2, 15], *Wrightia tinctoria* (Jain and Bari, 2010) [12]. Thus, Tribals in Pachamalai hills have a good knowledge about the use of

many plants for wounds and the plants prescribed by tribal healers are either based on single plant part or a combination of several plant parts. This preliminary effort helped to identify 26 species of commonly used plants for wounds. Further, studies have to be carried out to identify the numerous medicinal plants and the most important thing would be to conserve these plants. Dissemination of the knowledge of medicinal property would improve the socioeconomic status of the Malayali tribe.

References

- Adeleye OA, Femi-Oyewo MN, Bamiro OA *et al.* Ethnomedicinal herbs in African traditional medicine with potential activity for the prevention, treatment, and management of coronavirus disease 2019, Future Journal of Pharmaceutical Sciences. 2021;7:72.
- Babu G, Sanjeeva Bairy KI. Effect of *Tridax procumbens* on burn wound healing. International Journal of Advance Pharma and Bio Sciences 2003;40:488-491.
- Bhat RS, Shankrappa J, Shivakumar HG. Formulation and evaluation of polyherbal wound treatments, Asian Journal of Pharmaceutical Sciences. 2007;2:11-17.
- Cox PA, Ballick MJ. The ethnobotanical approach to drug discovery. Scientific American 1994;6:82-87.
- Fabricant DS, Farnsworth NR. The value of plants used in traditional medicine for drug discovery, Environmental Health Perspectives. 2001;109:69-75.
- Farnsworth NR, Soejarto DD. Potential consequences of plant extinction in the United States on the current and future availability of prescription drugs, Economic Botany. 1985;39(3):231-240.
- Gamble JS, Fischer CEC. Flora of the Presidency of Madras. Adlard & Sons Ltd., London, 1915-1936, 1-3,
- Gemechu W, Woldekidan S, Teka F *et al.* Ethnomedicinal uses, phytochemistry and pharmacological activities of *Rumex nervosus*, Journal of Analytical Pharmacy Research. 2021;10(2):65-69.
- Henry AN, Chithra V, Balakrishnan NP. Flora of Tamil Nadu. Botanical survey of India, Coimbatore 1989;1(3),
- Henry AN, Kumari GR, Chithra V. Flora of Tamil Nadu. Botanical Survey of India, Coimbatore. 1987;1:2.
- Ignacimuthu S, Ayyanar M. Ethnobotanical investigations among tribes in Madurai district of Tamil Nadu (India), Journal of Ethnobiology and Ethnomedicine. 2006;2(1):25.
- Jain PS, Bari SB. Evaluation of wound healing effect of petroleum ether and methanolic extract of *Abelmoschus manihot* (L.) Medik, Malvaceae and *Wrightia tinctoria* R. Br. Apocynaceae in rats, Revista Brasileira de Farmacognosia. 2010;20:756-61.
- Jain SK, Rao RR. A Hand book of Field and Herbarium Methods, Today and Tomorrow Printers and Publishers, New Delhi, 1976.
- Jain SK. Glimpses of Indian Ethnobotany. Oxford & IBH Publishing Co., New Delhi, 1981.
- Johnson DB, Gorle A. Evaluation of Ethanolic Extract of *Tridax procumbens* Linn on Wound Healing in Rats, Research Journal of Pharmacy and Technology. 2012;5(2):239-242.
- Mahapatra AD, Bhowmik P, Banerjee A, *et al.* Ethnomedicinal Wisdom: An Approach for Antiviral Drug Development. In (ed.) Ahmad Khan, M.S., Iqbal Ahmad and Chattopadhyay, D New Look to Phyto-medicine. 2019, 35-61.
- Matthew KM. Flora of the Tamil Nadu Carnatic. Rapinat

- Herbarium, Tiruchirappalli, 1981-1988, 1-3.
18. Nair NC, Henry AN. Flora of Tamil Nadu. Botanical Survey of India, Coimbatore. 1983;1:1.
 19. Pieroni A, Price LL, Vandebroek I. Welcome to Journal of Ethnobiology and Ethnomedicine. Journal of Ethnobiology and Ethnomedicine. 2005;1:1.
 20. Udupa SL, Udupa AL, Kulkarni DR. Influence of *Tridax procumbens* on dead space wound healing Fitoterapia. 1991;62(2):146-150.
 21. WHO, Traditional medicine Fact sheet no 1.3.4, 2003, <http://www.who.int/media/infocentre/factsheets/fs134/en/>.
 22. Williams LAD. Ethnomedicine. West Indian Medicine Journal. 2006;55(4):215.