



ISSN (E): 2320-3862  
ISSN (P): 2394-0530  
[www.plantsjournal.com](http://www.plantsjournal.com)  
JMPS 2022; 10(1): 125-135  
© 2022 JMPS  
Received: 19-11-2021  
Accepted: 22-12-2021

**Jiby John Mathew**  
Department of Biotechnology,  
Mar Augusthinose College,  
Ramapuram, Kerala, India

**Aswathy Nair**  
Department of Biotechnology,  
Mar Augusthinose College,  
Ramapuram, Kerala, India

**Sajeshkumar NK**  
Department of Biotechnology,  
Mar Augusthinose College,  
Ramapuram, Kerala, India

**Prem Jose Vazhacharickal**  
Department of Biotechnology,  
Mar Augusthinose College,  
Ramapuram, Kerala, India

**Corresponding Author:**  
**Prem Jose Vazhacharickal**  
Department of Biotechnology,  
Mar Augusthinose College,  
Ramapuram, Kerala, India

## Analysis of the phylloremediation capability of *Mangifera indica* in hydrocarbon polluted area: An outlook study

**Jiby John Mathew, Aswathy Nair, Sajeshkumar NK and Prem Jose Vazhacharickal**

### Abstract

Environmental pollution is reaching such great heights that it is a matter of global concern now. A phenomenon called phylloremediation was observed in plants, especially the ones that grow in polluted areas. Usually, plants growing in polluted areas themselves develop mechanisms to reduce the harmful pollutants in the surrounding to less harmful substances. This is done by the leaves and the microbes associated with the leaves either on their surface or inside the leaves. This project aims to analyze the phylloremediation capability in collected leaf samples from *Mangifera indica* growing in hydrocarbon polluted areas and to understand the various microbes associated with the leaves which are responsible for the degradation of air pollutants, the mechanisms involved and the genes or gene clusters which are responsible for this phenomenon. The procedure includes isolation of hydrocarbon-degrading bacteria from leaf samples, followed by its DNA and plasmid isolations, sequencing, and comparison with known gene sequences in databases using BLAST analysis. The whole process could be further enhanced by genetically engineering the selected microbe. Various genes that are already found to produce certain enzymes which can degrade one or more pollutants can be introduced into our selected organism's genome to improve the phylloremediation capability.

**Keywords:** Phylloremediation, *Mangifera indica*, hydrocarbon degraders, gene clusters, blast

### 1. Introduction

The contamination of the air by harmful or poisonous substances such that it greatly affects the health of all the living organisms on earth and at the same time harms the abiotic components of the ecosystem was referred to as air pollution. Environmental pollution is reaching such great heights that it is a matter of global concern now. The major reasons for such alarmingly increasing pollution rates are industrialization, increasing population, economic development, etc. The harmful effects of air pollution are not limited to health alone. It also affects the agriculture and economy which in turn have a great impact on human life.

Some of the major air pollutants are sulphur dioxide, nitrogen dioxides, carbon monoxide, ground-level ozone, particulate matter, and volatile organic compounds (Wei *et al.*, 2017) [47]. There has been a lot of effort being put into by various sectors of the society in trying to minimize the air pollution, but still, it seems to be just increasing day by day. Hence, we direct our attention to the remediation of air pollutants by exploiting the capabilities of plants, their different parts, and microorganisms associated with the plants.

Air pollutants can be mitigated through biological means, i.e., bioremediation. Bioremediation is defined as the process whereby toxic substances are biologically degraded under controlled conditions to non-toxic limits established by regulatory authorities (Mueller *et al.*, 1996) [28]. For effective bioremediation, enzymatic action of the microorganisms must take place. The major limitation of bioremediation is that it can take place only if the environmental conditions are suitable for microbial growth. Thus, there is a need for the manipulation of environmental parameters to allow microbial growth and degradation to proceed at a faster rate (Sharma, 2012) [44].

We are all familiar with phytoremediation. Phytoremediation is the use of plants to clean up soil, water, and in some cases, air, polluted with hazardous contaminants. The term is a combination of the Greek word-Phyto (plant) and the Latin word remedium (restoring balance).

It can be mainly employed for the removal of contaminants such as polycyclic aromatic hydrocarbons (PAHs), munition waste such as TNT, salt (NaCl), etc. Although it is a proven technology, its global usage as of now is very much limited. This is mainly because although theoretically, it might seem efficient, in reality, there hasn't been able to redress any of the environmental issues to a significant extent (Gerhardt *et al.*, 2017).

Phytoremediation is mainly employed in the clearing of soil and stagnant water bodies that are contaminated with substances like heavy metals (Cadmium, Lead, Arsenic, etc) (Feng *et al.*, 2013) [15]. Despite its encouraging results in the past years, many challenges have been encountered in phytoremediation in the field.

Phytoremediation is increasingly adopted in the treatment of soil and water pollution. However significant advances in the efficiency of the method are to be made for making phytoremediation a viable, economical option for pollution control.

All the above processes of phytoremediation are meant to reduce or eliminate mostly pollutants or contaminants in water and soil. In other words, phytoremediation mainly concentrates on the reduction of water pollution and soil pollution. In later researches, a different form of phytoremediation, called phylloremediation was observed in plants, especially the ones that grow in polluted areas. Usually, plants growing in polluted areas themselves develop mechanisms to reduce the harmful pollutants in the surrounding to less harmful substances. This is done by the leaves and the microbes associated with the leaves either on their surface or inside the leaves. Both leaves and the microbes can individually control air pollution but their combined effect enhances the process. Thus, phylloremediation is meant to reduce air pollutants whereas phytoremediation could deal with water and soil contaminants.

Phylloremediation is defined as “a natural process of bioremediation of air pollutants through leaves and leaf-associated microbes, not the microbes or leaves alone.” (Wei *et al.*, 2017) [47].

During the last three decades, air pollution is increasing due to economic growth and urbanization. Various policies and practices have been brought for the remediation of this pollution, but the air is still polluted. In this review, we are finding a way to reduce pollution using the potential of plant leaves and leaf-associated microbes. Plant leaves are capable of absorbing air pollutants or adsorbing them onto their surfaces and microbes on leaves are capable of converting the pollutants into non-toxic molecules (Wei *et al.*, 2017) [47].

## 2. Review of literature

A variety of microbes are found in the various parts of the plants. They include bacteria, fungi, yeast, algae, protozoa, nematodes. These microbes can either be present inside the leaf tissues or on the surface of leaves. Phyllosphere is the aerial habitat colonized by microbes. Though all types of microbes are found on the leaves, the majority of them are bacteria. Leaves provide a vast surface area for microbes. The surface area provided by leaves alone for microbes is around  $6.4 \times 10^8$  km sq. (Lindow and Brandl, 2003) [8].

Epiphytic bacterial populations differ sharply in size among and within plants of the same species, as well as nearby. These variations in population sizes are due to large fluctuations in the physical and nutritional conditions of the phyllosphere (Lindow and Brandl, 2003) [8].

### 2.1 Alteration of plant surface properties

Bacteria can produce compounds with surfactant properties. Hence, they increase the wettability of leaves (Neu *et al.*, 1990; Hutchison *et al.*, 1993; Bunster *et al.*, 1989) [29, 23, 9]. The cuticle is hydrophobic. Thus, the increased wettability of the leaf surface helps improve the solubility and diffusion of substrates. These can prove to be beneficial to the epiphytes. The bacterial movement on the surface of the leaves is also facilitated. The watery layer on the leaf surface produced by these surfactants helps the bacteria to move to places where nutrients are abundant.

Another factor commonly observed in some species which are mostly found as epiphytes are the production of the toxin, syringomycin by *Pseudomonas syringae* pv. *syringae* (Quigley *et al.*, 1994) [33]. This toxin induces the formation of ion channels. The plant metabolites are released and it leads to cell lysis (Hutchison *et al.*, 1995) [24]. Nonpathogenic strains of *P. syringae* produce syringomycin at levels that won't cause cell death but are enough to trigger the release of plant metabolites (Hutchison *et al.*, 1995) [24].

Many of the bacterial colonisers of the phyllosphere can produce indole-3-acetic acid, which is a plant growth regulator (Brandl *et al.*, 2001) [8]. This is of great importance as the bacteria colonising the leaves produce growth regulators essential for the plants and which are similar to the auxins produced by plants themselves (Lindow *et al.*, 2003) [8].

### 2.2 Development of phylloremediation technologies

So, as mentioned above, it is established that plants and microbes can individually contribute to the minimization of air pollutants, but their combined effort can accelerate this process. A series of steps have been introduced to develop phylloremediation technology which is as follows:

1. Plant selection
2. Microbe selection
3. Evaluation and compatibility between plant leaves and microbes
4. Implementation (Wei *et al.*, 2017) [47]

In their work, Wei and others selected plants from four categories: trees, shrubs, ground cover plants for use in outdoor environments, and foliage plants for outdoor environments.

Each plant responds to pollutants differently. Their tolerance levels vary and this is evaluated using the air pollution tolerance index (Singh *et al.*, 1991) [45]. Plants from each category were evaluated to check their tolerance against pollutants like PMs, O<sub>3</sub>, SO<sub>2</sub>, NO<sub>x</sub>, and VOCs and their capability to assimilate these (Wei *et al.*, 2017) [47]. Evaluation is done by exposing the seedlings to various pollutants at different concentrations and the analysis is made on factors like stomatal conductance, photosynthetic rate, morphological analysis, the maximum quantum efficiency of photosystem II, etc. (Wei *et al.*, 2017) [47].

After analysis, suitable plants with greater abilities to assimilate or absorb or adsorb pollutants and those with greater tolerance levels are selected. Thereafter, a suitable microbe is to be selected which is compatible with the selected plant. Some bacteria reside on the surface of the leaves and some others, called the endophytes, reside inside the plant tissues without harming the plants. Since endophytes are not washed away easily by any forms of precipitation, these are more desirable (Wei *et al.*, 2017) [47].

The whole process could be further enhanced by genetically engineering the selected microbe. Various genes that are

already found to produce certain enzymes which can degrade one or more pollutants can be introduced into our selected organism's genome to improve the phylloremediation capability. An example of this is the work done by Barac and others. When appropriate degradation pathways were introduced into the endophytic bacteria, they showed improved degradation of toluene. They introduced the pTOM toluene-degradation plasmid of *Burkholderia cepacia* G4 into *B. cepacia* L.S.2.4, a natural endophyte of yellow lupine. The engineered endophytic bacteria strongly degraded toluene (Barac *et al.*, 2004) [6].

Once the microbes are selected, they could be domesticated by exposing them to varying pH, carbon source, temperature, and oxygen. Their phylogenetic relationships with other microbes should be determined. Other factors to be determined and evaluated in the laboratory include their utilization of organic compounds, the decomposition rate of pollutants, adaptability, competition, and growth rate (Wei *et al.*, 2017) [47].

The major step following the selection and evaluation of both the plants and microbes in the evaluation of the compatibility between them. The microbes are inoculated into the selected plants to see if they can grow well on the leaf surfaces and at the same time to check if the inoculation affects plant growth (Wei *et al.*, 2017) [47]. Once compatible combinations are obtained, they are further tested by exposure to pollutants. The desired microbes can be propagated using bioreactors and plants by tissue culture or cuttings. For real-world testing, these plants could be transplanted to outside polluted environments or greenhouses (Wei *et al.*, 2017) [47].

### 2.3 Hydrocarbon pollution

Any compound which consists of carbon and hydrogen atoms is termed a hydrocarbon. They are organic compounds and some of the forms are gasoline, petroleum, coal, kerosene, charcoal, natural gas, etc. The majority of the air pollution is caused by the automobiles that run on these hydrocarbon-based fuels. It is the incomplete combustion of these hydrocarbon fuels that causes pollution. Hydrocarbons react with oxides of nitrogen to form gases which when are hit by the sun's rays leads to the formation of ozone, which is a heavy form of oxygen and settles down in the atmosphere. Hence, the polluted air feels heavy. The incomplete combustion of hydrocarbons thus affects the quality of air in several ways and causes severe health issues.

### 2.4 Hydrocarbon degrading bacteria

Hydrocarbons are present on the surface of plant parts, either as by-products or end products of metabolism. Leaves have a waxy coating that contains hydrocarbons (Ilori *et al.*, 2006). Microorganisms with the capability to oxidize various hydrocarbons are widely distributed. The presence of hydrocarbon on leaves attracts more hydrocarbon degraders to colonize it. Thus, there is an increase in the populations of hydrocarbon degraders. These microbes end up on these surfaces by the action of rain, wind, vector, or brush contact (Atlas, 1978) [5]. Thus, such phylloplane act as training sessions for such hydrocarbon-degrading bacteria (Ilori *et al.*, 2006).

The large number of hydrocarbon degraders on the leaves studied indicate that these organisms have been exposed to hydrocarbons well in advance, which would be mainly from the leaves. Organisms, however, reach the leaf surfaces through dust, air current, and rain splashes. The presence of nutrients such as exudates or waxes from the cuticle is the

driving factor for the establishment of these microbes on the new phylloplane (Rao, 1977) [34].

The seven different genera of hydrocarbon degraders isolated from the different plants that Ilori and others studied were: *Acinetobacter*, *Flavobacterium*, *Bacillus*, *Pseudomonas*, *Alcaligenes*, *Corynebacterium*, and *Micrococcus* (Ilori *et al.*, 2006).

Out of these, the ones that were isolated from *Mangifera indica* and thus found to be compatible with the phylloplane of *Mangifera indica* were: *Pseudomonas* sp., *Alcaligenes* sp., and *Micrococcus* sp. (Wei *et al.*, 2017) [47].

### 2.5 *Pseudomonas citronellois*

*Pseudomonas citronellois* can degrade hydrocarbons such as isoprenoid compounds including citronellol (Seubert *et al.*, 1960) [43]. And hence it got its present name. It was also found to be able to degrade complex oily sludge contaminations (Bhattacharya *et al.*, 2003) [7]. Thus, the species might have the potential to serve as bioremediation organisms to clean up oil-contaminated sites.

The strain P3B5 was isolated from plant leaf material. Plant leaves are covered by a cuticle. This cuticle consists of an esterified aliphatic polymer called cutin. It is overlaid by intra- and epi-cuticular waxes consisting of very-long-chain aliphatic compounds, including alkanes, alcohols, and fatty acids (Remus-Emsermann *et al.*, 2016) [35]. *P. citronellois* can degrade aliphatic compounds too. Thus, there is a possibility that the bacterium may be able to exploit aliphatic compounds as a nutrient source during leaf surface colonization (Remus-Emsermann *et al.*, 2016) [35]. Furthermore, *P. citronellois* were also found to have the ability to degrade terpenoid compounds (Forster-Fromme and Jendrossek, 2006) [16].

Two Alk-B-like alkane monooxygenase genes are present in the genome of *Pseudomonas citronellois*, which might have the potential to nutritionally access long and very-long-chain alkanes available on plant leaves (Remus-Emsermann *et al.*, 2016) [35].

### Alkane 1-Monooxygenase 1

Gene sequence which encodes the protein Alkane 1-monooxygenase 1 is given below

```
ATGAAAGCGCTGCATTATCTGAAATATTTTCTGTTTC
ATGCGGTGGCCTGCTGAGCGCG
GCGGCGCTGCTGGCGGGCGGCAGCTGGATTGCGTGC
GGCTGGCGGCGGTGCTGACCATT
TATGTGCTGGGCGATCTGGTGTGCGGCGATGATACC
AGCACCCCGGAATTTTCATCATCCG
GGCATTCTGACCTTTCAGCTGTGGCTGGCGCTGCCG
TGCTGTGCTTTATTGTGTTTGGC
GCGCTGTGGGGCGTGAGCGATGGCGATTTTCTGGGC
TTTGGCGGCGCTGGGCCAGGCG
ACCGGCTATGATCTGATTGCGGCGCGATGCGACC
AGCCTGGGCCAGCATGTGTGCGCG
TGGATTCTGACCGGCTGGTATTGGCACCATTGGC
ACCATTACCGCGCATGAACTGACC
CATCGCACCTGGGATCCGGTGAGCATGCTGATTGGC
CGCTGGCTGCTGGCGTTTAGCTTT
GATACCATTTTTAGCATTGAACATGTGTATGGCCATC
ATCGCTATGTGAGCACCACCGAA
GATCCGGCGACCGCGCCGCGCGGCAACGTGTAT
GCGCATGTGCTGATTAGCACCGTG
CGCGCAACCTGAGCGCGTGGCATATTGAAAAACAG
CGCCTGGCGCGCAAAGGCTATTGG
CTGCTGGGCTGGCGCAACGCGGTGCTGCGGCCAT
```

GGCATGAGCCTGCTGCTGGTGGCG  
 GCGGCGTATGCGATTGGCCAGTGGCCGGCGGCGCTG  
 TTTTTTATTGCGTGC GCGCTGTGG  
 GGCAAAGCGCTGCTGGAAATTGTGAACTATATGGAA  
 CATTATGGCATGGTGC GCAACCCG  
 GCGACCCCGGTGCAGCCGCGCCATAGCTGGAACACC  
 AACAAACGCATTAGCAGCTGGACC  
 ATGTTTAACTGACCCGCCATAGCCATCATCATGCG  
 CAGGGCGAAGTGCCGTATCAGGAT  
 CTGAAACCGTTTCCGGATGCGCCGATGATGGTGGGC  
 GGCTATCTGACCACCATATTGTG  
 GCGATGCTGCCGCGCTGTGGCATCGCCTGATGACC  
 CCGAAAGTGCTGGCGTGGGATCGC  
 GAATATGCGAGCGCGGAAGAACGCCAGCTGGCGGC  
 GACCGCGAACGCGCGCAGCGGCATT  
 CCGGCGCTGATGAAATATAGCAACTATCCGAGCCCC  
 AGC

(Retrieved from  
[https://www.bioinformatics.org/sms2/rev\\_trans.html](https://www.bioinformatics.org/sms2/rev_trans.html))

The fasta format of the protein Alkane 1- monooxygenase 1 is given below

MKALHYLKYFLFHAVGLLSAAALLAGGSWIACGLAA  
 VLTIVVLGDLVCGDDTSTPEFHHPGILTFQLWLA  
 LPLLCFIVFAALWGVSDGDFLGFGAALGQATGYDLIAA  
 RDATSLGQHVC AWILTGLVIGTIGTITAHFLT  
 HRTWDPVSMILGRWLLAFSFDITFSIEHVYGHHRVST  
 TEDPATAPRGRNVYAHVLISTVRGNLSAWHIE  
 KQRLARKGYWLLGWRNAVLRGHGMSLLLVAAYAI  
 GQWPAALFFIACALWGKALLEIVNYMEHYGMVRNP  
 ATPVQPRHSWNTNKRISSWTMFLNRHSHHHAQGEVP  
 YQDLKPFDPAPMMVGGYLTIIIVAMLPLWHRL  
 MTPKVLAWDREYASAEERQLAATANARSGIPALMKY  
 SNYPSPS

(NCBI, accession no. AMO76609)

### Alkane 1-Monooxygenase 2

Gene sequence which encodes the protein Alkane 1- monooxygenase 2 is given below:

ATGTTTTCGTTTCTGAGCCCGCGTGGATGCTGCGCC  
 TGA AAAAAGCGGGCTATTGGATT  
 TGGCTGGTGCCGGTGTGGCATTCCGCTGAGCTATT  
 GGTGGAGCTATGGCAGCGGCCAT  
 CCGAACCGGTGGGCGTGGTTTGTGATTGCGGTGGTG  
 TTTCTGGTGATTCCGGTGCTGGAT  
 GCGATTGTGGGCCGCGATCCGGCGAACCCGGATGAA  
 GCGAGCGAAGTGCCGGCGCTGGAA  
 CGCGAAGGCTATTATCGTTTTCTGAGCCTGATGACC  
 GTGCCGCTGCTGCTGGGCATGCTG  
 GTGTATGGCGGCTGGGTGATTACCCATTATGAAGCG  
 TGGA ACTGGGTGGGCAA ACTGGGC  
 TGGACCCTGAGCGTGGGCACCGTGATGGGCGCGATT  
 GGCATTACCGTGAGCCATGAACTG  
 ATTCATAAAGATCCGCAGCTGGAACAGCGCGCGGGC  
 GGCCTGCTGCTGGCGGCGGTGTGC  
 TATGCGGGCTTTAAAGTGGAACATGTGCGCGGCCAT  
 CATGTGCATGTGAGCACCCCGGAA  
 GATGCGAGCAGCAGCCGCTATGATCAGAGCCTGTAT  
 GCGTTTCTGCCGCATGCGTATAAA  
 CATAACTTTCTGAACGCGTGGCGCCTGGAAGCGCAG  
 CGCCTGAAACGCAAAGGCCTGCCG  
 GCGCTGCATTGGCGCAACGA ACTGATTTGGTGGTAT  
 GCGGTGAGCGCGCTGTTTCTGTCTG  
 GGCTTTAGCTGCGCTTTGGCTGGCTGGGCGCGGTG  
 TATTTTCTGGGCCAGGCGCGATG  
 GCGTTTACCCTGCTGGAAATTGTGAACTATGTGGAA

CATTATGGCCTGCATCGCCGCCCGC  
 CTGGATGATGGCCGCTATGAACGCACCACCCATGAA  
 CATAGCTGGAACAGCAACTTTCTG  
 CTGACCAACCTGTTTCTGTTTCATCTGCAGCGCCATA  
 GCGATCATCATGCGTATGCGAAA  
 CGCCGCTATCAGGTGCTGCGCCATTATGATGATAGC  
 CCGCAGCTGCCGAACGGCTATGCG  
 GGCATGATTGTGCTGGCGCTGTTTCCGCCGCTGTGGC  
 GCGCGGTGATGAACCCGCGCGTG  
 CGCGCGTATTATGCGGGCGAAGAATATCAGCTGAGC  
 GATACCCAGCGCGCG

(Retrieved from  
[https://www.bioinformatics.org/sms2/rev\\_trans.html](https://www.bioinformatics.org/sms2/rev_trans.html))

The fasta format of the protein Alkane 1- monooxygenase 2 is given below

MFAFLSPAWMLRLKKAGYWIWLVFVFIPLSYWWSY  
 GSGHPNAWAWFVIAVFLVIPVLDIAVGRDPANP  
 DEASEVPALEREGYRFLSLMTVPLLLGMLVYGGWVI  
 THYEAWN WVGKLGWTLVSGTVMGAIGITVSHL  
 IHKDPQLEQRAGGLLLAAVCYAGFKVEHVRGHHVHS  
 TPEDASSRYDQSLYAFPHAYKHNFLNAWRLE  
 AQRLKRKGLPALHWRNELIWWYAVSALFLLGFSLAFG  
 WLGAVYFLGQAAMAFTLLEIVNYVEHYGLHRRR  
 LDDGRYERTTHEHSWNSNFLTNLFLFHLQRHSDHHA  
 YAKRRYQVLRHYDDSPQLPNGYAGMIVLALFPP  
 LWRAVMNPRVRAYYAGEEYQLSDTQRA  
 (NCBI, accession no. AMO75838)

*P. citronellolis* P3B5 contains two putative AlmA-like monooxygenases responsible for the degradation of very-long-chain alkanes (>C30) (Remus-Emsermann *et al.*, 2016) [35].

### FAD-Containing Monooxygenase EthA

Gene sequence which encodes this protein is given below

ATGAGCGTGGAACATCTGGATGTGCTGATTGTGGGC  
 GCGGGCCTGAGCGGCATTGGCGCG  
 GCGTATCATCTGATGAAACATTGCCCGGGCAAACCC  
 TTTGCGCTGCTGGAAGGCCGCGAA  
 GCGATGGGCGGCACCTGGGACTGTGTTTCGCTATCCG  
 GGCATTTCGAGCGATAGCGATATG  
 TATACCCTGGGCTATAACTTTAAACCGTGACCGAT  
 GCGAAAGCGATTGCGGATGGCCCCG  
 AGCATTTCGCCATTATATTGAAGAAACCGCGCAGGAA  
 AACGGCATTGATCGCAA AATTCGC  
 TATCGCCATCGCGTGCTGAAAGCGGATTGGAACAGC  
 GATAGCGCGCGCTGGCTGCTGACC  
 GTGCAGCGCGCGCATGAACCGGAACCGGTGCGCAT  
 GAGCGCGCAGTTTCTGCTGATGTGC  
 ACCGGCTATTATCGCTATGAAGCGGGCTATAACCCCG  
 GAATTTAAAGGCCGCGATAGCTTT  
 CGCGGCCAGATTATTCATCCGCAGCTGTGGCCGGAA  
 GGCTTTGATTATGCGGATAAACCG  
 GTGGTGGTATTGGCAGCGGCGGACCGCGGTGACC  
 CTGGTGCCGAGCCTGACCGATAAA  
 GCGCGCCATGTGACCATGCTGCAGCGCAGCCCGAGC  
 TATGTGATTAGCTGCCGAGAAA  
 GATCCGATTAGCAACTTTCTGCGCCGCTTTCTGCCGG  
 AAACCTGGGTGTATCGCCAGGCG  
 CGCGCGCGCAACGTGACCATGCAGATGGTGTTTTTT  
 CTGTTTGCGAAAACCTTTCCGAGC  
 CTGGCGCGCAAAGCGCTGCTGGGCGCTGGCGCGCCAT  
 CAGCTGGGCAAAGATTTTGATATG  
 CGCCATTTAGCCCGCGCTATAAACCGTGGGATGAA  
 CGCGTGTGCTGCGTGCCGATGGC  
 GATCTGTTTAAAGCGCTGCGCAAAGGCAAAGCGAGC

GTGGTGACCGAACATATTGATGGC  
 TTTTGCGAAAAAGGCATTCGCCTGAAAAGCGGCCAG  
 GTGCTGGAAGCGGATGTGATTGTG  
 ACCGCGACCGGCCTGGATCTGGTGATGTTTGGCGGC  
 GCGGAACCTGGCGGTGGATGGCAA  
 CCGTTTCAGGTGAACCAGAGCATGGGCTATCGCGGC  
 ATTATGCTGCGCGATCTGCCGAAC  
 CTGGCGGGCGGTGGTGGGCTATACCAACGCGAGCTGG  
 ACCCTGAAAGCGGATCTGAGCAGC  
 GAATATTTTTGCCGCCTGATTAACCATATGGATGCGA  
 TTGGCATGCGCCAGTGCACCCCG  
 CGCGCGGGCGCGGGCCAGGTGAAAGAAGAACCCTT  
 TCTGAACCTGAACAGCGGCTATATT  
 CAGCGCGCGGCGGATAAAATGCCGAAACAGGGCGA  
 TCGCACCCCGTGGAAACTGTATCAG  
 AACTATGCGCTGGATCTGGCGCTGCTGCCCATGGC  
 AAATGGAAGATGCTATCTGACC  
 TTTACCAGCCCGCGCGCCAGGGCGATGAAGCG  
 AGCCTGCAGGCGGTG

([https://www.bioinformatics.org/sms2/rev\\_trans.html](https://www.bioinformatics.org/sms2/rev_trans.html))

The fasta format of the protein is given below:

MSVEHLVDLIVGAGLSGIGAAAYHLMKHCPGKTFALLE  
 GREAMGGTWDLFRYPGIRSDSDMYTLGYNFKPW  
 TDAKAIADGPSIRHYIEETAQENGIDRKIRYRHRVLKAD  
 WNSDSARWLLTVQRGDEPEPVRMSAQFLLMC  
 TGYRYEAGYTPEFKGRDSFRGQIIHPQLWPEGFDYAD  
 KRVVVIGSGATAVTLVPSLTDKARHVTMLQRS  
 PSYVISLPQKDPISNFLRRFLPETWVYRQARARNVTMQ  
 MVFFLFAKTFPSLARKALLGLARHQLGKDFDM  
 RHFSRPYKPWDERVCCVPDGLFKALRKGKASVVTEH  
 IDGFCEKGIRLKSQVLEADVIVTATGLDLVMF  
 GGAEALVDGKPFQVNSMGYRIGIMLRDLPNLAADVVG  
 YTNASWTLKADLSSEYFCRLINHMDAIGMRQCTP  
 RAGAGQVKEEPFLNLSGYIQRAADKMPKQGDRTWP  
 KLYQNYALDLALLRHGKVEDGYLFTSPRARQGD  
 EASLQAV

(NCBI, accession no. AMO73862)

*P. citronellolis* is able to degrade terpenes such as citronellol. A corresponding gene cluster (PcP3B5\_19950–20010) was also detected in *P. citronellolis* P3B5, which may be responsible for enabling this species to colonize the phyllosphere (Remus-Emsermann *et al.*, 2016) [35].

*P. citronellolis* P3B5 genome contains genes whose products are involved in the biosynthesis of indole-3-acetic acid (PcP3B5\_05210–05220, PcP3B5\_17810, PcP3B5\_07120–07140). It is a plant hormone of the auxin class and induces plant cell elongation and division, which leads to an increase in locally available nutrients. Thus *P. citronellolis* P3B5 has an impact on the plant host (Remus-Emsermann *et al.*, 2016) [35].

## 2.6 Pseudomonas putida

*Pseudomonas putida* has been found to colonize the phyllosphere of several plants and its various strains are involved in the active degradation of various hydrocarbons. One of them is the hydrocarbon- naphthalene. Various strains of *Pseudomonas putida* contain certain plasmids that contain genes that lead to the degradation of naphthalene.

## 2.7 Naphthalene

Naphthalene is the simplest polycyclic aromatic hydrocarbon with the formula C<sub>10</sub>H<sub>8</sub>. It is a white crystalline solid with a characteristic odour, detectable at concentrations as low as

0.08 ppm by mass (Amoore *et al.*, 1983) [4]. Its structure consists of a fused pair of benzene rings. Exposure to large amounts of naphthalene may damage or destroy red blood cells (Santucci and Shah, 2000) [36], cause confusion, nausea, vomiting, diarrhea, blood in the urine, and jaundice. The International Agency for Research on Cancer (IARC) classifies naphthalene as possibly carcinogenic to humans and animals. The IARC also points out that acute exposure causes cataracts in humans, rats, rabbits, and mice; and that hemolytic anaemia can occur in children and infants after oral or inhalation exposure or after maternal exposure during pregnancy.

## 2.8 Pseudomonas putida G7

*Pseudomonas putida* G7 contains the NAH7 naphthalene catabolic plasmid. The nah operons are divided into two clusters (Undugoda *et al.*, 2016) [46]. The genes of the upper operon have genes encoding enzymes that convert naphthalene to salicylate. The lower operon has genes encoding enzymes that convert salicylate to acetyl coenzyme A and pyruvate (Zhou *et al.*, 2002) [49].

A Tn5 insertion mutation which occurs immediately upstream from the first gene of the lower operon, nahG, found in the NAH7 plasmid gives the Nah<sup>-</sup> Sal<sup>-</sup> phenotype (Schell, 1985; Park *et al.*, 2002) [39], which defined the regulatory region containing the gene encoding the NahR protein. The NahR protein is a LysR-type transcriptional regulator (LTTR). This protein is necessary for activation of both the upper (genes nahA-F) and lower (genes nahG-M) operons on the NAH7 plasmid (Huang & Schell, 1991; Schell, 1985, 1986, 1993; Schell *et al.*, 1990; Schell & Poser, 1989; Schell & Wender, 1986; Park *et al.*, 2002) [22, 39, 40, 41, 42, 37, 38]. Studies have shown that NahR is constitutively expressed at low levels (Cebolla *et al.*, 1997) [11]. NahR, bound to its inducer, salicylate, leads to the activation of both nah operons by interacting with DNA enhancer sequences upstream of the promoter region in upper and lower nah operons (Schell & Poser, 1989) [37]. The 5' flanking region of known nahR genes is the divergently transcribed nahG-nahR promoter region and NahR binding in this -60 bp region upstream of the nahG gene transcription start site induces nahG gene transcription, but represses its own expression by negative autoregulation (Schell & Poser, 1989; Schell & Wender, 1986; Yen & Serdar, 1988) [37, 38, 48]. The exact mechanism of autoregulation is yet to be discovered (Yen & Serdar, 1988) [48].

## 2.9 Pseudomonas putida NCIB 9816

*P. putida* NCIB 9816 is capable of utilizing naphthalene as the sole carbon source. An 81 kb plasmid, pDTG1 containing nahR gene conferred this trait in this strain. The plasmid encodes key enzymes in early enzymatic steps in naphthalene degradation (Undugoda *et al.*, 2016) [46].

A few of the gene segments present on the plasmid pDTG1 and the proteins coded by them are mentioned below:

### nahR Gene

The Gene sequence of the nahR gene found in plasmid pDTG1 in *Pseudomonas putida* NCIB 9816-4 is given below  
 ATGGAACCTGCGTGATCTGGATTTAAACCTGCTGGTG  
 GTGTTCAACGAGCTGCTGGTCTGCAGACCGCTCT  
 CTATCGTTGCCGAGAACCTGGCCCTGCCAGCCAGCCTG  
 CTGTCAGCAATGCGTGAAACGCCTGCGCACAGC  
 GCTACAGGACCAACTTTTCGTGCGCACCTACCAGGG  
 TATGGAACCCACACCCTACGCCGGAACCTTGCC

GAGCCCGTCGCTTGGCCATGCACGCCCTGCGCGAA  
 GCCCTGCAGACCATGAGCGCTTCGATCCGCTGA  
 CCAGCGAGCGTACTTTCACCCTGGCCATGACCGACA  
 TCGGCGAGATCTACTTCATGCCGCGGCTGATGGA  
 TGCGATTACTCTTCAAGCCCCCAATTGCGCGATCAG  
 CACCGTGCAGGACAGTTCGATGAGCCTGATGCAA  
 GCCTTGACAGACGGCACAGTGGACCTAGCCGTGGGC  
 CTGCTGCCAACCTGCAGACTGGCTTCTTCCAGC  
 GCCGGTGTCTCCACAATCACTACGTGTGCTGTGTC  
 GCAAGGACCATCCGGCCACCCGCGAACCCCTGAC  
 TCTTGAGCGCTTCTGTTTCTATGGCCACGTGCGTGTG  
 ATCGCCGCTGGCACAGGCCACGGCGAGGTGGAC  
 ACATACTTGACGAAGGCCGGCATCCGGCGCGACATC  
 CGCTGGAGGTGCCCACTTCGTGCGCCGTTGGCC  
 ACATACTCCAGCGCACCGAGCTGCTGCCACTGTGC  
 CGATACTGTTTCGCCGACTGCTGCTAGAGCCCTT  
 CGGCTGAGCGTCTTGGCCACCCAGTGCCTTGGCC  
 GGAAATCGCCATCAACATGTTTGGCATGCGAAA  
 TACCACCAGGACCTCGCCAATATCTGGTTGCGGCAA  
 CTGATGTTTCGACCTGTTTTCTGATTGA

(Retrieved from NCBI; Accession ID: AF491307)

The protein encoded by this gene is a naphthalene degradation LysR-family transcriptional activator.

MELRDLNLLVFNELLVCRVSIVAENLGLTQPAVS  
 NALKRLRTALQDQLFVRTYQMEPTPYAANLA  
 EPVALAMHALREALQHHERFDPLTERTFTLAMTDIGE  
 IYFMPRLMDAITLQAPNCAISTVRDSSMSLMQ  
 ALQNGTVDLAVGLLPNLQTFQRRLLHNHYVCLCRK  
 DHPATREPLTLERFCSYGHVRIAAGTGHGEVD  
 TYLTKAGIRRDIREVPHFVAVGHILQRTELLATVPIRF  
 ADCCVEPFLSVLPHVALPEIAINMFWHAK  
 YHQDLANIWLRQLMFDLFS  
 (NCBI; Accession ID: AAM09539)

## 2.10 Naphthalene dioxygenase enzyme

The first step in the aerobic degradation of naphthalene is catalyzed by naphthalene dioxygenase (NDO) enzyme system (EC 1.14.12.12) from *Pseudomonas* sp. strain NCIB 9816-4. In this reaction, both atoms of oxygen are added to the aromatic nucleus of naphthalene by NDO, forming cis-naphthalene dihydrodiol (Jeffrey *et al.*, 1975; Jerina *et al.*, 1971) [25, 26]. The NDO system consists of three components, each of which has been purified and characterized (Rebecca *et al.*, 2000). An iron-sulfur flavoprotein reductase and an iron-sulfur ferredoxin, both, transfer electrons from NAD(P)H to the catalytic oxygenase component (Ensley and Gibson, 1983; Ensley *et al.*, 1982; Haigler and Gibson, 1990) [14, 13, 19]. The oxygenase is composed of a large subunit called  $\alpha$  and a small subunit called  $\beta$ , in an  $\alpha\beta\beta$  configuration (Kauppi *et al.*, 1998) [27]. Naphthalene dioxygenase is a member of a large family of oxygenases whose  $\alpha$  subunits contain a Rieske [2Fe-2S] center and mononuclear nonheme iron (Butler and Mason, 1997). Electrons are transferred from the Rieske center of the ferredoxin to the Rieske center of the oxygenase  $\alpha$  subunit. The reduced Rieske center in one  $\alpha$  subunit transfers an electron to mononuclear iron at the active site in an adjacent  $\alpha$  subunit (Kauppi *et al.*, 1998; Parales *et al.*, 1999) [27, 31]. His-208, His-213, and Asp-362 coordinate the active-site iron, forming a 2-His-1-carboxylate facial triad. Asp-205 in the catalytic domain of the NDO  $\alpha$  subunit is hydrogen-bonded to His-208 and His-104 in the adjacent  $\alpha$  subunit. His-104 is one of the Rieske center ligands. Asp-205 is required for efficient electron transfer from the Rieske center to the active-site iron (Parales *et al.*, 1999) [31].

## *P. putida* naphthalene dioxygenase gene (nahAa, nahAb, nahAc, and nahAd)

### nahAb gene

ATGGAACCTCATACAACCGAACAAATCGCATAATT  
 CCCTTCAGTGCCGGTGCCAACCTTCTGGAAGTGC  
 TTCGCGAGAACGGTGTAGCTATTTCTACAGTTGCTT  
 GTCTGGGCGTTGCGGAACCTGTGCTGCCGGGT  
 TATAGATGGCAGTGTGATTGATTCTGGGGCGGAAAA  
 TGGGCAATCAAACCTCACCGACAAGCAGTATGTG  
 CTCGCCGTGTCAGTCACTACTGGAATTGCGCTA  
 TCGAAGTCCCAGAAGCCGACGAAATTGTCACTC  
 ACCCGGCGCAATCATCAAGGGCACAGTGGTTCGCG  
 TCGAGTCCGCACTCACGATATCCGTCGCTTACG  
 CGTACGCCTCTCCAAGCCCTTCGAGTTCTACCCGGA  
 CAGTACGCGACACTGCAGTTCAGCCCTGAGCAT  
 GCGCGTCCGATTCATGAGCAGTTCGCGAGTACG  
 CAAGAAATGGAGTTCACATACGCAAGTGGCCGG  
 GTGGGCGGTCACGGAGTATGTTTTCGAACACGTCC  
 GCGAAGGTACAAGCATCAAGTTGAGCGGCCTCT  
 TGGTACGGCTTATCTACGTCAGAAGCACACCGGACC  
 GATGCTGTGTGATTGTTGCGGGGACCGGACTCGCA  
 CCGGTGCTGTCGATTGTTGCGGGCGGCTGAAGTGC  
 GGTATGACGAACCCATCCTCCTTTATTTTCGGGG  
 TGCGCAGTCAAGACCTCTACGACGACAGAGCGAT  
 TGCACAACTCGCCGTGACCACCTCAACTGAC  
 CGTACACACGGTATTGCAACGGGCCCGATTAATGA  
 GGGTACGCGAGCCGGCCTAATTACCGATGTGATC  
 GAAAAGACATCCTTTTCGCTGGCTGGGTGGAGGGCC  
 TACCTGTGCGGCGCACCGAGCGATGGTTGAAGCGT  
 TGTGCACCGTCACCAAGCATCTTGAATATCACCCG  
 AACATATTTATGCCGATGCCTTCTATCCCGGTGG  
 GATCTGA

(NCBI; Accession ID: AF491307 REGION: 14536.15522)

## It encodes naphthalene 1, 2-dioxygenase reductase component (plasmid)

(*Pseudomonas putida*), whose FASTA sequence is as follows:

MELLIQPNRIIPFSAGANLLEVLRENGVAISYSCLSGR  
 CGTCRCRVIDGSVIDSGAENGQSNLTDKQYV  
 LACQSVLTGNCAIEVPEADEIVTHPARIKGTVVAVESP  
 THDIRRLRVRLSKPFEFSPGYATLQFSPEH  
 ARPYSMAGLPDDQEMEFHIRKVPGGRVTEYVFEHVRE  
 GTSIKLSGPLGTAYLRQKHTGPMLCVGGGTGLA  
 PVLIVRGALKSGMTPILLYFGVRSQQDLYDAERLHK  
 LAADHPQLTVHTVIATGPINERAGLITDVI  
 EKDILSLAGWRAYLCGAPAMVEALCTVTKHLGISPEHI  
 YADAFYPGGI

(NCBI; Accession ID: AAA25904)

The purified enzyme catalyzed the reduction of cytochrome c, dichlorophenolindophenol, Nitro Blue Tetrazolium, and ferricyanide. These activities were enhanced in the presence of added FAD. The ability of the enzyme to catalyze the reduction of the ferredoxin involved in naphthalene reduction and other electron acceptors indicates that it functions as a NAD(P)H-oxidoreductase in the naphthalene dioxygenase system. The results suggest that naphthalene dioxygenase requires two proteins with three redox groups to transfer electrons from NADH to the terminal oxygenase (Haigler and Gibson, 1990) [19].

### nahAb gene

ATGCAGTAAAGTGGATTGAAGCAGTCGCTCTTTCT  
 GACATCCTTGAAGGTGACGTCCTCGGCGTGACTG  
 TCGAGGGCAAGGAGCTGGCGCTGTATGAAGTTGAAG

CGCAAATCTACGCTACCGACAACCTGTGCACGCA  
TGGTCCGCCCGCATGAGTGATGGTTATCTCGAGGG  
TAGAGAAATCGAATGCCCTTGCATCAAGGTCGG  
TTTGACGTTTGCACAGGCAAAGCCCTGTGCGCACCC  
GTGACACAGAACATCAAACATATCCAGTCAAGA  
TTGAGAACCTGCGCGTAATGATTGATTTGAGCTAA  
(NCBI. Accession ID: AF491307 REGION: 15666..15980)  
It encodes naphthalene 1, 2-dioxygenase ferredoxin  
component (plasmid) (*Pseudomonas putida*), whose FASTA  
sequence is as follows:  
MTVKWIEAVALSDILEGDVLTVEGKELALYEVEGEI  
YATDNLCTHGSARMSDGYLEGREIECPLHQGR  
FDVCTGKALCAPVTQNIKTYPVKIENLRVMIDLS  
(NCBI; Accession ID: AAA25905)

One of the three components of the naphthalene dioxygenase,  
the protein contained 2 g-atoms each of iron and acid-labile  
sulphur. It had an apparent molecular weight of 13,600. The  
evidence suggests that it is a ferredoxin-type protein that  
functions as an intermediate electron transfer protein in  
naphthalene dioxygenase activity (Haigler and Gibson, 1990)  
[19].

#### NahAc gene

ATGAATTACAATAATAAAATCTTGGTAAGTGAATCT  
GGTCTGAGCCAAAAGCACCTGATTCATGGCGATG  
AAGAACTTTTCCAACATGAACTGAAAACCATTTTTG  
CGCGGAACTGGCTTTTTCTCACTCATGATAGCCT  
GATTCCTGCCCCGGCGACTATGTTACCGCAAAAAT  
GGGATTGACGAGGTCATCGTCTCCCGGCAGAAC  
GACGGTTCGATTCGTGCTTTTCTGAACGTTTGCCGGC  
ATCGTGGCAAGACGCTGGTGAGCGTGGAAGCCG  
GCAATGCCAAAGGTTTTGTTTGCAGCTATCACGGCT  
GGGGCTTCGGCTCCAACGGTGAAGTGCAGAGCGT  
TCCATTTGAAAAAGATCTGTACGGCGAGTCGCTCAA  
TAAAAATGTCTGGGGTTGAAAGAAGTCGCTCGC  
GTGGAGAGCTTCCATGGCTTCATCTACGGTTGCTTCG  
ACCAGGAGGCCCTCTCTTATGGACTATCTGG  
GTGACGCTGCTTGGTACCTGGAACCTATGTTCAAGC  
ATTCCGGCGGTTTAGAACTGGTTCGCTCCCTCCAGG  
CAAGGTTGTGATCAAGGCCAACTGGAAGGCACCCCG  
GGAAAACCTTTGTGGGAGATGCATACCACGTGGGT  
TGGACGACGCGTCTTCGCTTCGCTCGGGGGAGTCT  
ATCTTCTCGTTCGCTCGCTGGCAATGCGGCGCTAC  
CACCTGAAGGCGCAGGCTTGC AAATGACCTCCAAAT  
ACGGCAGCGGCATGGGTGTGTTGTGGGACGGATA  
TTCAGGTGTGCATAGCGCAGACTTGGTTCGGGAATT  
GATGGCATTTCGAGGCGCAAAGCAGGAAAGGCTG  
AACAAAGAAATTGGCGATGTTTCGCGCTCGGATTTAT  
CGCAGCCACCTCAACTGCACCGTTTTCCCGAACA  
ACAGCATGCTGACCTGCTCGGGTGTGTTTCAAAGTAT  
GGAACCCGATCGACGCAAACACCACCGAGGTCTG  
GACCTACGCCATTGTGCAAAAAGACATGCCTGAGGA  
TCTCAAGCGCCGCTTGGCCGACTCTGTTACGCGA  
ACGTTTCGGGCTGCTGGCTTCTGGGAAAGCGACGAC  
AATGACAATATGGAAACAGCTTCGCAAAAACGGCA  
AGAAATATCAATCAAGAGATAGTGATCTGCTTTCAA  
ACCTTGGTTTCGGTGAGGACGTATACGGCGACGC  
GGTCTATCCAGGCGTTCGTCGGCAAATCGGGCAGTCGG  
CGAGACCAGTTATCGTGGTTTCTACCGGGCTTAC  
CAGGCACACGTCAGCAGCTCCAAGTGGGCTGAGTTC  
GAGCATGCCTCTAGTACTTGGCATACTGAACTTA  
CGAAGACTACTGATCGCTAA  
(NCBI; Accession ID: AF491307 REGION: 16051..17400)  
It encodes naphthalene 1,2-dioxygenase iron sulfur protein  
component large subunit (plasmid) (*Pseudomonas putida*),

whose FASTA sequence is as follows:  
MNYNNKILVSEGLSQKHLIHGDEELFQHELKTIFARN  
WLFLTHDSLIPAGDYVTAKMGIDEVIVSRQN  
DGSIRAFNLVCRHRGKTLVSVEAGNAKGFVCSYHGW  
GFGSNGELQSVPEKDLYGESLNKKCLGLKEVAR  
VESFHGFIYGCDFQEAPPLMDYLGDAAWYLEPMFKHS  
GGLELVGPPGKVVIKANWKAPAENFVGDAYHVG  
WTHASSLRSGESIFSSLAGNAALPEAGLQMTSKYGS  
GMGVLWDGYSVHVSADLVPPELMAFGGAKQERL  
NKEIGDVRARIYRSHLNCTVFPNNSMLTCSGVFKVWN  
PIDANTTEVWTYAIVEKDMPELKRRLADSVQR  
TFGPAGFWESDDNDNMETASQNGKQYQSRSDLLSNL  
GFGEDVYGDVYPGVVGKSAIGETSIRGFYRAY  
QAHVSSSNWAEFEHASSTWHELTKTTDR  
(NCBI; Accession ID: AAO64274)

#### NahAd gene

ATGATGATCAATATTCAAGAAGACAAGCTGGTTTCC  
GCCCACGACGCCGAAGAGATTCTTCGTTTCTTCA  
ATTGCCACGACTCTGCTTTGCAACAAGAAGCCACTA  
CGCTGCTGACCCAGGAAGCGCATTTGTTGGACAT  
TCAGGCTTACCGTGCTTGGTTAGAGCACTGCGTGGG  
GTCAGAGGTGCAATATCAGGTCATTTACGCGAA  
CTGCGCGCAGCTTCAGAGCGTCGTTATAAGCTCAAT  
GAAGCCATGAACGTTTACAACGAAAATTTTCAGC  
AACTGAAAGTTCGAGTTGAGCATCAACTGGATCCGC  
AAAAGTGGGGCAACAGCCCCGAAGCTGCGCTTTAC  
TCGCTTTATCACCAACGTCCAGGCCGCAATGGACGT  
AAATGACAAAGAGCTACTTCACATCCGCTCCAAC  
GTCATTCTGCACCGGGCAGCAGCTGGCAATCAGGTC  
GATGTCTTCTACGCCGCCCGGGAAGATAAATGGA  
AACGTGGCGAAGGTGGAGTACGAAAATTTGGTCCAGC  
GATTCGTCGATTACCCAGAGCGCATACTTCAGAC  
GCACAATCTGATGGTCTTTCTGTGA  
(Source: NCBI; Accession ID: AF491307 REGION:  
17415..17999)

It encodes naphthalene 1,2-dioxygenase iron sulfur protein  
component small subunit (plasmid) (*Pseudomonas putida*),  
whose FASTA sequence is as follows:

MMINIQEDKLVSAHDAEEILRFNCHDSALQQEATLL  
TQEAHLDDIQAYRAWLEHCVGSEVQYQVISRE  
LRAASERRYKLNEAMNVYNENFQQLKVRVEHQLDPQ  
NWGNSPKLRFTRFITNVQAAMDVNDKELLHIRSN  
VILHRARRGNQVDVFYAAREDKWKRGEVVRKLVQR  
FVDYPERILQTHNLMVFL  
(Source: NCBI; Accession ID: AAO64275)

### 3. Materials and Methods

#### 3.1 Sample collection

Collect the leaves of *Mangifera indica* from the streets where  
vehicular traffic was high. The leaves were transported to the  
lab on a sealed cover and stored in the refrigerator until use.

#### 3.2 Isolation of total heterotrophic and hydrocarbon utilizing bacteria from leaf samples

Each leaf sample (4 g) should be washed with 100 ml of  
phosphate buffer and then shaken at 200 rev/min for one hour.  
Then the diluted samples should directly be added to the  
modified mineral salt agar plates. Plates are to be then  
incubated at room temperature (28°C-30°C) for five days.  
Bacterial colonies should then be streaked on PAH added  
Bacto-Bushnell Haas medium to select PAH utilizing bacteria  
(Undugoda *et al.*, 2016) [46].

#### 3.3 Characterization of the isolates

Each isolate should be examined for its size, shape, margin,

consistency, opacity, elevation, pigmentation, Gram reaction, and cell morphology as described by Cowan (1974) [12]. The isolates are to be characterized as described by Holt *et al.* (1994) [21]. Diagnostic properties used should include motility, production of cytochrome oxidase, catalase, indole and urease, gelatin liquefaction, starch hydrolysis, oxidation/fermentation of sugars, methyl red test, Voges Proskauer test, and growth at 42°C and 5°C (Ilori *et al.*, 2006).

### 3.4 Characterization of the isolates

Each isolate should be examined for its size, shape, margin, consistency, opacity, elevation, pigmentation, Gram reaction, and cell morphology as described by Cowan (1974) [12]. The isolates are to be characterized as described by Holt *et al.* (1994) [21]. Diagnostic properties used should include motility, production of cytochrome oxidase, catalase, indole and urease, gelatin liquefaction, starch hydrolysis, oxidation/fermentation of sugars, methyl red test, Voges Proskauer test, and growth at 42°C and 5°C (Ilori *et al.*, 2006).

### 3.5 Selection of efficient PAH degrading bacteria

The best PAH degrading bacterial strains can be selected based on the results obtained from the colorimetric and HPLC methods indicated below:

#### Colorimetric assay

Each bacterial strain is inoculated into Bacto Bushnell-Haas broth incorporated with PAH compound (1% v/v) and Methylene blue (2% v/v), the redox indicator and incubated at room temperature (28°C-30°C) with constant shaking at 180 rev/min, for 14 days with control without bacterial inoculation. From broth culture 5 ml sample is centrifuged at 6000 rev/min for five minutes. The recovered supernatant is assayed spectrophotometrically by measuring absorbance at 609 nm for the residual hydrocarbon. Six replicates are done for each bacterial strain and PAH degradation percentage is determined using the following equation (Okafor *et al.*, 2009) [30].

The absorbance of sample Percentage of PAH degradation =  $1 - (\text{Absorbance of sample} / \text{Absorbance of Control}) \times 100$

#### HPLC determination of PAH degradation

Each bacterial strain is inoculated into Bacto Bushnell-Haas broth incorporated with PAH (phenanthrene, naphthalene, etc.) compound (100 ppm). Then it is incubated at room temperature (28°C-30°C) with constant shaking at 180 rev/min, for 14 days with control without bacterial inoculation.

Residual PAH in the culture is extracted with hexane and acetone-containing mixture. Then extract is analyzed by high-performance liquid chromatography (HPLC) equipped with a UV detector. The analytical column (250 mm long, 4.6 mm diameter) is packed porous spherical C-18 material (packed size, 5µm). Acetonitrile-water mixture (75: 25) is used as a mobile phase for PAHs at a flow rate of 1.0 ml min<sup>-1</sup>. The sample (20 µL) is injected into the column through the sample loop. UV - detector is set at 254 nm for compound detection. The Chromleon chromatography software is used for the quantification of PAHs throughout the experiment. Finally, the percentage of degradation is determined (Undugoda *et al.*, 2016) [46].

### 3.6 Isolation of catabolic plasmid and confirmation of plasmid-encoded PAH degradation

Catabolic plasmids of the isolated microbes can be isolated according to the method of (Sambrook *et al.*, 1989) and each plasmid is to be transformed into the E.coli JM109. PAH degrading ability of transformants are to be tested using

colorimetric and HPLC method to determine the plasmid based PAH degradation. Plasmid based PAH degradation is confirmed by curing the plasmid of bacterial cells using acridine orange (Fujii *et al.*, 1997) [17] and then testing their PAH degradation ability by colorimetric and HPLC methods (Undugoda *et al.*, 2016) [46].

### 3.7 PCR amplification

Suitable primers are designed for the PCR amplification of the required genes. Plasmid DNA templates are amplified using the above primers. Then PCR products are visualized and their size estimation is done by gel electrophoresis (Undugoda *et al.*, 2016) [46].

### 3.8 Sequencing of PCR products

Automated sequencing should be carried out using applied Biosystems automated sequencer (ABI3730XL). Purified PCR products representing the gene segments are sequenced directly using appropriate primers (Undugoda *et al.*, 2016) [46].

### 3.9 Restriction digestion of PCR products

PCR amplicons of the genes can be digested using the Hind III restriction enzyme to determine their RFLP patterns (Undugoda *et al.*, 2016) [46].

### 3.10 BLAST analysis

Nucleotide sequences of the gene segments are to be aligned with gene sequences of Genbank using BLAST (Altschul *et al.*, 1997) [3].

## 4. Expected results and discussion

The outcomes expected on the completion of the described methodology are.

1. Many species of microorganisms were expected in the *Mangifera* leaves.
2. There is a chance for hydrocarbon utilizing microorganisms in the isolate.
3. Isolation of DNA from the selected organisms.
4. DNA Sequences.
5. BLAST result.

If the genome sequence obtained after the sequencing, matches with any of the bacterial sequences mentioned earlier, it can be able to concluded that the leaves of *Mangifera indica* collected also have one or more of these microbes associated with it and it was developed as a result of heavy hydrocarbon pollution in the area that it was growing in.

And hence it could be said that the plant selected also performs phylloremediation. On the other hand, if the sequences of the microbes isolated from the leaf samples don't match with the gene sequences of the organisms that were earlier found out to perform phylloremediation, it is highly unlikely that the plant selected has phylloremediation capabilities.


## 5. Conclusions

Environmental pollution is reaching such great heights that it is a matter of global concern now. Out of all the measures being taken to reduce this menace, a very cheap and effective approach could be - the use of the concept of phylloremediation of air pollutants, as the aerial surfaces of leaves are estimated to sum up to 4x10<sup>8</sup> sq. km. on the earth (Wei *et al.*, 2017) [47]. Objective of this study was to analyse the phylloremediation capability in collected leaf samples from *Mangifera indica* growing in hydrocarbon polluted areas and to understand the various microbes associated with the




leaves which are actually responsible for the degradation of air pollutants, the mechanisms involved and the genes or gene clusters which are responsible for this phenomenon. This information can definitely be used later to induce this natural capability of plants and microbes into the ones that lack it

through genetic engineering, followed by tissue culture and micropropagation to produce a large number of genetically modified plants that'll be able to degrade various air pollutants.



Source: [https://media.springernature.com/lw785/springer-static/image/chp%3A10.1007%2F978-3-540-85467-8\\_3/MediaObjects/978-3-540-85467-8\\_3\\_Fig1\\_HTML.jpg](https://media.springernature.com/lw785/springer-static/image/chp%3A10.1007%2F978-3-540-85467-8_3/MediaObjects/978-3-540-85467-8_3_Fig1_HTML.jpg)

Fig 1: NAH7 plasmid of *Pseudomonas putida* strain G7.



Source: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC94462/>

Fig 2: Reaction catalyzed by the three-component NDO system.

**6. Acknowledgment:** The authors are grateful for the cooperation of the management of Mar Augustinose College for necessary support. Technical support from Binoy A Mulantra is also acknowledged.

**Conflicts of interest:** The author do not have any conflict of interest.

**7. References**

1. Abeshu MA, Geleta B. Medicinal uses of honey. *Biology and Medicine*. 2016; 8(2):1.
2. Ahangari Z, Naseri M, Vatandoost F. Propolis: chemical composition and its applications in endodontics. *Iranian Endodontic Journal*. 2018;13(3):285.
3. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, et al. Gapped BLAST and PSI-BLAST: a new

- generation of protein database search programs. *Nucleic Acids Res.* 1997;25:3389-3402.
4. Amoore JE, Hautala E. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatiles for 214 industrial chemicals in air and water dilution. *J Appl Toxicology.* 1983;3(6):272-290. doi:10.1002/jat.2550030603. PMID 6376602. S2CID 36525625.
  5. Atlas RM. Stimulated petroleum degradation. *Crit. Rev. Microbiol.* 1978;5:371-386.
  6. Barac T, Taghavi S, Borremans B, Povoost A, Oeyen L, Colpaert JV *et al.* Engineered endophytic bacteria improve phytoremediation of water-soluble, volatile, organic pollutants. *Nat. Biotechnol.* 2004;22:583-588. doi: 10.1038/nbt960
  7. Bhattacharya D, Sarma PM, Krishnan S, Mishra S, Lal B. Evaluation of Genetic Diversity among *Pseudomonas citronellolis* Strains Isolated from Oily Sludge-Contaminated Sites. *Appl Environ Microbiol.* 2003;69:1435-41.
  8. Brandl MT, Quinones B, Lindow SE. Heterogeneous transcription of an indoleacetic acid biosynthetic gene in *Erwinia herbicola* on plant surfaces. *Proc. Natl. Acad. Sci. USA.* 2001;98:3454-3459.
  9. Bunster L, Fokkema HJ, Schippers B. Effect of surface activity of *Pseudomonas* spp. on leaf wettability. *Appl. Environ. Microbiol.* 1989;55:1340-1345.
  10. Butler CS, Mason JR. Structure-function analysis of the bacterial aromatic ring-hydroxylating dioxygenases. *Adv Microb Physiol.* 1997;38:47-84.
  11. Cebolla A, Sousa C, de Lorenzo V. Effector specificity mutants of the transcriptional activator NahR of naphthalene degrading *Pseudomonas* define protein sites involved in binding of aromatic inducers. *J Biol Chem.* 1997;272:3986-3992.
  12. Cowan ST. Cowan and Steel's Manual for the identification of medical bacteria. Second edition. Cambridge University Press, 1974.
  13. Ensley BD, Gibson DT, Laborde AL. Oxidation of naphthalene by a multicomponent enzyme system from *Pseudomonas* sp. strain NCIB 9816. *J Bacteriol.* 1982;149:948-954.
  14. Ensley BD, Gibson DT. Naphthalene dioxygenase: purification and properties of a terminal oxygenase component. *J Bacteriol.* 1983;155:505-511.
  15. Feng R, Wei C, Tu S. The roles of selenium in protecting plants against abiotic stresses. *Environmental and Experimental Botany.* 2013;87:58-68. doi:10.1016/j.envexpbot.2012.09.002.
  16. Forster-Fromme K, Jendrossek D. Identification and characterization of the acyclic terpene utilization gene cluster of *Pseudomonas citronellolis*. *FEMS Microbiol Lett.* 2006;264:220-5.
  17. Fujii T, Takeo M, Maeda Y. Plasmid encoded genes specifying aniline oxidation from *Acinetobacter* sp. strain YAA. *Microbiology.* 1997;143:93-99.
  18. Gerhardt KE, Gerwing PD, Greenberg BM. Opinion: Taking phytoremediation from proven technology to accepted practice, Dec-2016. Retrieved from <https://pubmed.ncbi.nlm.nih.gov/28167031/>
  19. Haigler BE, Gibson DT. Purification and properties of ferredoxin NAP, a component of naphthalene dioxygenase from *Pseudomonas* sp. strain NCIB 9816. *J Bacteriol.* 1990;172:465-468.
  20. Haigler BE, Gibson DT. Purification and properties of NADH-ferredoxin NAP reductase, a component of naphthalene dioxygenase from *Pseudomonas* sp. strain NCIB 9816. *J Bacteriol.* 1990;172:457-464.
  21. Holt JG, Krieg NR, Sneath PHA, Staley JT, Williams ST. *Bergey's Manual of Determinative Bacteriol.* Williams and Wilkins Baltimore, 1994.
  22. Huang, JZ, Schell MA. *In vivo* interactions of the NahR transcriptional activator with its target sequences. Inducer-mediated changes resulting in transcription activation. *J Biol Chem.* 1991;266:10830-10838.
  23. Hutchison ML, Johnstone K. Evidence for the involvement of the surface active properties of the extracellular toxin tolaasin in the manifestation of brown blotch disease symptoms by *Pseudomonas tolaasii* on *Agaricus bisporus*. *Physiol. Mol. Plant Pathol.* 1993;42:373-384.
  24. Hutchison ML, Tester MA, Gross DC. Role of biosurfactant and ion channel-forming activities of syringomycin in transmembrane ion flux: a model for the mechanism of action in the plant-pathogen interaction. *Mol. Plant-Microbe Interact.* 1995;8:610-620.
  25. Jeffrey AM, Yeh HJC, Jerina DM, Patel TR, Davey JF, Gibson DT *et al.* Initial reactions in the oxidation of naphthalene by *Pseudomonas putida*. *Biochemistry.* 1975;14:575-583.
  26. Jerina DM, Daly JW, Jeffrey AM, Gibson DT. cis-1, 2-Dihydroxy-1, 2-dihydronaphthalene: a bacterial metabolite from naphthalene. *Arch Biochem Biophys.* 1971;142:394-396.
  27. Kauppi B, Lee K, Carredano E, Parales RE, Gibson DT, Eklund H *et al.* Structure of an aromatic ring-hydroxylating dioxygenase-naphthalene 1,2-dioxygenase. *Structure.* 1998;6:571-586.
  28. Mueller JG, Cerniglia CE, Pritchard PH. "Bioremediation of environments contaminated by polycyclic aromatic hydrocarbons," in *Bioremediation: Principles and Applications*, eds E. L. Crawford, D. L. Crawford (Cambridge: Cambridge University Press), 1996, 125-194.
  29. Neu TR, Nartner T, Poralla K. Surface active properties of viscosin: a peptide lipid antibiotic. *Appl. Microbiol. Biotechnol.* 1990;32:518-520.
  30. Okafor UG, Tasié F, Okafor FM. Hydrocarbon degradation potential of indigenous fungal isolated from petroleum contaminated soils. *The journal of physical and natural science.* 2009;3:56-68
  31. Parales RE, Parales JV, Gibson DT. Aspartate 205 in the catalytic domain of naphthalene dioxygenase is essential for activity. *J Bacteriol.* 1999;181:1831-1837.
  32. Parales RE, Lee K, Resnick SM, Jiang H, Lessner DJ, Gibson DT *et al.* Substrate specificity of naphthalene dioxygenase: effect of specific amino acids at the active site of the enzyme. *Journal of bacteriology.* 2000;182(6):1641-1649. <https://doi.org/10.1128/JB.182.6.1641-1649.2000>
  33. Quigley NB, Gross DC. Syringomycin production among strains of *Pseudomonas syringae* pv. *syringae*: conservation of the syrB and syrD genes and activation of phytotoxin production by plant signal molecules. *Mol. Plant-Microbe Interact.* 1994;7:78-90.
  34. Rao SNS. Microflora of plant surface especially leaf surface and biochemical reactions. In: *Soil microorganism and Plant Growth.* Oxford and IHB. Publishing Company, 1977, 75-80.
  35. Remus-Emsermann *et al.* Complete genome sequence of

- Pseudomonas citronellolis* P3B5, a candidate for microbial phyllo-remediation of hydrocarbon-contaminated sites, 2016. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5037603/>
36. Santucci K, Shah B. Association of naphthalene with acute hemolytic anemia. *Acad Emerg Med.* Jan-2000;7(1):42-7. doi:10.1111/j.1553-2712.2000.tb01889.x. PMID 10894241
  37. Schell MA, Poser EF. Demonstration, characterization, and mutational analysis of NahR protein binding to nah and sal promoters. *J Bacteriol.* 1989;171:837-846.
  38. Schell MA, Wender PE. Identification of the nahR gene product and nucleotide sequences required for its activation of the sal operon. *J Bacteriol.* 1986;166:9-14.
  39. Schell MA. Transcriptional control of the nah and sal hydrocarbon-degradation operons by the nahR gene product. *Gene.* 1985;36:301-309.
  40. Schell MA. Homology between nucleotide sequences of promoter regions of nah and sal operons of NAH7 plasmid of *Pseudomonas putida*. *Proc Natl Acad Sci USA.* 1986;83:369-373.
  41. Schell MA. Molecular biology of the LysR family of transcriptional regulators. *Annu Rev Microbiol.* 1993;47:597-626.
  42. Schell MA, Brown PH, Raju S. Use of saturation mutagenesis to localize probable functional domains in the NahR protein, a LysR-type transcription activator. *J Biol Chem.* 1990;265:3844-3850.
  43. Seubert W. Degradation of isoprenoid compounds by microorganisms I., *Pseudomonas citronellolis* n. sp: Isolation and Characterization of an Isoprenoid-Degrading Bacterium. *J Bacteriol.* 1960;79:426-34.
  44. Sharma S. Bioremediation: Features, Strategies and applications. *Asian Journal of Pharmacy and Life Science,* 2012, 2(2). Retrieved from <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.1086.2035&rep=rep1&type=pdf>
  45. Singh SK, Rao DN, Agrawal M, Pandey J, Narayan D. Air pollution tolerance index of plants. *J. Environ. Manag.* 1991;32:45-55. doi: 10.1016/S0301-4797(05)80080-5
  46. Undugoda LJS, Kannangara S, Sirisena DM. Genetic Basis of Naphthalene and Phenanthrene Degradation by Phyllosphere Bacterial Strains *Alcaligenes faecalis* and *Alcaligenes* sp. 11SO. *J Bioremed Biodeg.* 2016;7:333. doi:10.4172/2155-6199.1000333
  47. Wei X, *et al.* Phylloremediation of Air Pollutants: Exploiting the Potential of Plant Leaves and Leaf-Associated Microbes, 2017. Retrieved from <https://www.frontiersin.org/articles/10.3389/fpls.2017.01318/full>
  48. Yen KM, Serdar CM. Genetics of naphthalene catabolism in pseudomonads. *Crit Rev Microbiol.* 1988;15:247-268.
  49. Zhou NY, Al-Dulayymi J, Baird MS, Williams PA. Salicylate 5-hydroxylase from *Ralstonia* sp. strain U2: a monooxygenase with close relationships to and shared electron transport proteins with naphthalene dioxygenase. *J Bacteriol.* 2002;184:1547-155.