



ISSN (E): 2320-3862
ISSN (P): 2394-0530
<https://www.plantsjournal.com>
JMPS 2023; 11(6): 70-72
© 2023 JMPS
Received: 03-08-2023
Accepted: 10-09-2023

Gurmanpreet Singh Sodhi
General Shivdev Singh Diwan
Gurbachan Singh Khalsa
College, Patiala, Punjab, India

Kamalesh Kumar
General Shivdev Singh Diwan
Gurbachan Singh Khalsa
College, Patiala, Punjab, India

Effect of growth and yield of maize (*Zea mays* L.) by optimization of nitrogen through organic and inorganic sources crop under climatic conditions of Punjab

Gurmanpreet Singh Sodhi and Kamalesh Kumar

Abstract

Experiment was conducted during *zaid* season of 2020-21 at the Agricultural Research Farm Dhablan of the G.S.S.D.G.S. Khalsa College Patiala, Punjab. Field experiment was laid out in split plot design with 16 different treatments with 3 replications, combination of organic and inorganic fertilizer levels. The soil of experimental field was clay in texture with pH 7.6 and organic carbon 0.58%, available nitrogen 184.4 kg ha⁻¹, available phosphorus 19.10 kg ha⁻¹ and available K 144.9 kg ha⁻¹. Different levels of nitrogen also influenced the plant height at 30, 60 and 90 days. Plant height frequently increases with increase in nitrogen levels during all stages of crop growth. Maximum plant height, number of leaves and dry weight with nitrogen application @ 125% ha⁻¹ whereas the lower plant height, number of leaves and dry weight found over the control treatment (N₀). Increase in height, number of leaves and dry weight might be due to positive effect of nitrogen on the growth of the plant which leads to rapid increase in length of internodes and plant height, number of leaves and dry weight. Maximum biological yield among all the nitrogen level, 217.91 q ha⁻¹ data were recorded with N_{125%} application which is significantly higher over the control (175.31 q ha⁻¹). Grain yield increased with FYM₂₀ application over the control which was 73.97 q ha⁻¹. High level of nitrogen increased the grain yield which was near about 72.97 q ha⁻¹. Similarly maximum harvest index (37.07%) was found at highest nitrogen level of N₀ while the minimum harvest index (33.43%) seen in control treated plots.

Keywords: Maize, plant height, seed yield

Introduction

Maize (*Zea mays* L.) is the third most versatile emerging food grain crop. In India, maize ranks third in terms of production among the cereals. It is widely adapted crop and is grown under extreme diverse climatic conditions ranging from tropical to temperate regions. There is no cereal in this world which has so immense potential and that is why it is also represented with other name as queen of cereals. Maize in America and India is referred to as corn which literally means "that which sustains life". Botanically, it belongs to grass family *Graminaceae/Poaceae*. Maize has tremendous variability in kernel color, texture, composition and appearance. It is an annual plant with an extensive fibrous root system. It is a diploid species, with a chromosome number of 2n = 20 and it followed C₄ pathway.

Maize is normally a tall annual grass with a solid stem. Usually in maize plant, 12-18 leaves are present in it. Being a monoecious plant it has two types of inflorescence i.e. female inflorescence which develops into an ear also known as cob and the male inflorescence which contains the male flowers also known as tassel. Generally, maize grain contains about 10% protein, 4% oil, 70% carbohydrate, 2.3% crude fiber, 1.4% ash, 10% albuminoids and substantial quantities of Vitamin A, Nicotinic acid, Riboflavin, Vitamin E (Singh *et al.* 2012) [8].

Materials and Methods

Experiment was conducted during *zaid* season of 2020-21 at the Agricultural Research Farm Dhablan is situated at about 24-46 °N latitude and 76-24 °E longitude at an altitude of about 250 m above the mean sea level. The experiment was laid out in split plot design with 3 replications. From the five randomly selected plants the heights were recorded in cm. The numbers of leaves were counted from the sample plants and the values of these were averaged.

Corresponding Author:
Gurmanpreet Singh Sodhi
General Shivdev Singh Diwan
Gurbachan Singh Khalsa
College, Patiala, Punjab, India

To study the dry weight of five plants were collected from the sampling rows of each plot at 30 days interval from sowing till harvest of the crop. These fresh samples were air dried and then dried in an oven at 60 °C till a constant weight was obtained and weighed to record the average dry weight of the plant. The weight of the sun dried harvested crop was recorded from net plot area and expressed in quintal per hectare after subtracting the seed yield. Seed yield of each plot excluding the border and sampling row was weighed in kilo gram and converted into $q\ ha^{-1}$.

Results and Discussion

In case of FYM, plant height was also affected significantly at all stages. Application of FYM @ 20 t ha^{-1} produced the tallest plant height (29.51 cm, 95.11 cm, 169.76 cm) at 30, 60 and 90 DAS over the control (FYM₀) which is (25.51 cm, 87 cm, 156.90 cm). Different levels of nitrogen also influenced the plant height at 30, 60 and 90 days. Plant height frequently increases with increase in nitrogen levels during all stages of crop growth. Highest plant height (31.55cm, 100.87cm and 174.72 cm) observed at 30, 60 and 90 DAS with nitrogen application @ 125% ha^{-1} whereas the lower plant height (23.11 cm, 78.29 cm and 147.09 cm) found over the control treatment (N₀). Increase in height might be due to positive effect of nitrogen on the growth of the plant which leads to rapid increase in length of internodes and plant height. Verma (2011) [10] research shows similar result as he identified that plant height increases with increase in the nitrogen levels.

In addition, a significant response seen in term of number of leaves with the application of FYM observed at all stages of observation which were (4.95 cm, 9.81 cm, 11.70 cm) but minimum number of leaves were observed in the control replication (3.82 cm, 8.18 cm and 10.29 cm) during 30 DAS interval. Analysis of data revealed that nitrogen had a significant effect on number of leaves plant⁻¹. More number of leaves were produced with optimum nitrogen dose as compared to low nitrogen doses at all stages. Highest number of leaves plant⁻¹ were recorded (5.62cm, 10.81cm and 12.90 cm) at N_{125%} ha^{-1} .

In FYM treated plots, FYM20t ha^{-1} accumulated high dry weight of plant which is near about 29.34 g plant⁻¹, 143.73 g plant⁻¹, 247.18 g plant⁻¹, 313.93 g plant⁻¹ at 30, 60, 90 and at harvest stage which is significantly higher over the control at all stages of crop growth. Among the nitrogen treated plots, N_{125%} produced higher dry matter (29.67 g, 147.99 g, 250.84 g and 331.21 g plant⁻¹) at 30, 60, 90 and at harvest stage over the low level of nitrogen treated plots which is N_{75%} and control. This is due to high photosynthetic rate in crop help to increase the total dry matter production of the plant. FYM treatment had a significant effect on grain yield. Maximum grain yield (73.97 q ha^{-1}) was found with FYM 20t ha^{-1} and minimum grain yield (62.4 q ha^{-1}) was found over the control. Nitrogen had also a significant effect on the yield of the grains. Maximum yield (72.97 q ha^{-1}) was found with the nitrogen application of the N_{125%} and minimum grain yield (65 q ha^{-1}) was found at the control plots. The reason might be due to increase in nitrogen level increases the photosynthetic rate and increase in the efficiency of the carbohydrates. Significant higher stover yield was found in the FYM 20t ha^{-1} which was 134.54 q ha^{-1} while 129.24 q ha^{-1} yield also obtained from the treatment FYM10t ha^{-1} which is much more over the control 122.87 q ha^{-1} . The higher stover yield also found by Sujatha, *et al.* 2008 [9] and Raman and Suganya

(2018) [13] through INM approach. Increasing in nitrogen levels also boost up the stover yield. Similarly maximum straw yield (144.94 q ha^{-1}) was observed under the nitrogen application of N_{125%} and minimum stover yield (110.31q ha^{-1}) was recorded at the control plots. FYM significant affects the biological yield also. Maximum biological yield (208.51 q ha^{-1}) was found with 20t ha^{-1} FYM treatment and minimum biological yield (185.27 q ha^{-1}) was found over the control. This is due to continuous supply of nutrient will increase the biological yield.

Maximum biological yield (217.91 q ha^{-1}) was found at high nitrogen dose N_{125%} and minimum biological yield (175.31 q ha^{-1}) was found at the control plots. This might be due to nitrogen level increases better crop growth rate and leaf area index which automatically enhances the biological yield. Similar results were found by Rajashekhar *et al.* 2019 [12] and Mishra *et al.* 2019 [1]. Both treatment i.e. FYM and Nitrogen shows significant effect on harvest index. The highest harvest index (36.23%) recorded with FYM10t ha^{-1} treated plots in comparison to FYM0, FYM 5t and FYM20t ha^{-1} . Similar results were found by Shah *et al.* 2017 [4] and Sharma *et al.* 2019 [6]. Similarly maximum harvest index (37.07%) was found at highest nitrogen level of N₀ while the minimum harvest index (33.43%) seen in control treated plots. Yigermal *et al.* 2019 [12] and Yadav *et al.* 2019 [11].

Table 1: Effect of farmyard manure and inorganic nitrogen level on periodic plant height of *zaid* maize

Treatments	Plant height (cm)			
	30DAS	60DAS	90 DAS	At harvest
Farmyard Manure (t ha^{-1})				
0 t FYM	23.01	84.58	154.73	163.37
5 t FYM	26.56	90.43	162.38	173.53
10 t FYM	28.62	93.38	166.61	177.44
20 t FYM	29.51	95.11	169.76	180.62
S.Em(±)	0.36	0.338	0.295	0.564
CD 5%	1.412	1.326	1.159	2.216
Nitrogen (% of recommended dose)				
0% N	22.03	78.29	147.34	161.55
75% N	26.39	88.62	162.54	172.05
100% N	28.48	96.56	169.57	178.22
125% N	30.8	100.04	174.05	183.14
S.Em(±)	0.656	0.568	1.519	1.35
CD 5%	1.914	1.657	4.435	3.941

Table 2: Effect of farmyard manure and inorganic nitrogen level on periodic Number of leaves plant⁻¹ of *zaid* maize

Treatments	Number of leaves plant ⁻¹			
	30DAS	60DAS	90 DAS	At harvest
Farmyard Manure (t ha^{-1})				
0 t FYM	3.82	8.18	10.29	9.59
5 t FYM	4.29	8.76	10.81	10.25
10 t FYM	4.64	9.31	11.28	10.72
20 t FYM	4.95	9.81	11.7	11.13
S.Em(±)	0.051	0.139	0.118	0.132
CD 5%	0.2	0.545	0.464	0.52
Nitrogen (% of recommended dose)				
0% N	3.2	7.39	9.41	8.78
75% N	3.95	8.34	10.3	9.82
100% N	4.94	9.52	11.47	10.93
125% N	5.62	10.81	12.9	12.17
S.Em(±)	0.163	0.238	0.137	0.359
CD 5%	0.476	0.694	0.401	1.047

Table 3: Effect of farmyard manure and inorganic nitrogen level on periodic dry weight of *zaid* maize

Treatments	Dry weight (g plant ⁻¹)			
	30DAS	60DAS	90 DAS	At harvest
Farmyard Manure (t ha⁻¹)				
0 t FYM	23.36	126.88	224.02	281.9
5 t FYM	27.97	135.78	237.51	294.87
10 t FYM	28.82	140.31	241.69	306.94
20 t FYM	29.34	143.73	247.18	313.93
S.Em(±)	0.259	0.447	0.309	0.283
CD 5%	1.016	1.756	1.212	1.111
Nitrogen (% of recommended dose)				
0% N	23.72	123.63	219.33	264.74
75% N	27.23	133.53	235.63	288.53
100% N	28.86	141.54	244.6	313.16
125% N	29.67	147.99	250.84	331.21
S.Em(±)	0.535	0.625	1.104	0.697
CD 5%	1.56	1.824	3.224	2.033

Table 4: Effect of farmyard manure and inorganic nitrogen level on grain yield, stover yield and biological yield of *zaid* maize

Treatments	Grain yield (q ha ⁻¹)	Stover yield (q ha ⁻¹)	Biological yield (q ha ⁻¹)	Harvest Index
Farmyard Manure (t ha⁻¹)				
0 t FYM	62.4	122.87	185.27	33.83
5 t FYM	67.68	126.18	193.86	35.02
10 t FYM	73.15	129.24	202.39	36.23
20 t FYM	73.97	134.54	208.51	35.55
S.Em(±)	0.342	0.304	0.537	0.103
CD 5%	1.342	1.195	2.108	0.403
Nitrogen (% of recommended dose)				
0% N	65	110.31	175.31	37.07
75% N	67.79	122.63	190.42	35.56
100% N	71.44	134.94	206.39	34.57
125% N	72.97	144.94	217.91	33.43
S.Em(±)	0.772	0.666	1.067	0.28
CD 5%	2.255	1.943	3.115	0.817

Conclusion

Among the FYM levels, the better and maximum output regarding crop growth and yield were seen in 20 t ha⁻¹ of FYM when it was applied during the initial stage of crop growth which helps to increase the soil physical, chemical and biological composition. However, maximum return seen in FYM₁₀ due to its lesser treatment cost than FYM₂₀ treatment. In-fact, in sub treatment, nitrogen plays an important role in terms of yield and growth attributes. N_{125%} shows statistically higher result.

References

- Mishra P, Tiwari US, Pandey HP, Pathak RK, Sachan AK. Impact of INM on growth and yield of maize (*Zea mays*) crop in central plain zone of Uttar Pradesh, India. International Journal of Current Microbiology and Applied Sciences. 2019;8(4):138-150.
- Rajashekhar L, Gurumurthy KT, Vageesh TS, Dhananjaya BC, Ganapathi, Sridhar CJ. Effect of integrated nutrient management practices on growth and yield of rainfed maize in sigehadlu micro watershed of Chikkamagaluru district. International Journal of Chemical Studies. 2019;6:58-61.
- Narang KK, Jha SK. Effect of weed management on weed dynamics, crop growth, fodder yield and economics of fodder maize (*Zea mays* L.) African tall under Chhattisgarh condition. Int. J Adv. Chem. Res. 2022;4(2):235-241.
DOI: 10.33545/26646781.2022.v4.i2d.104
- Shah RA, Wani BA. Yield, Nutrient Uptake and Soil Fertility of Maize (*Zea mays* L.) as Influenced by Varying Nutrient Management Practices under Temperate Conditions of Kashmir Valley, India. National Agriculture Innovation Project, Kumpwara; c2017. p. 75-78.
- Nuru Seid Tehulie. Review on critical period of weed competition and management in maize (*Zea mays* L.). Int. J Horticult Food Sci. 2021;3(2):44-48.
- Sharma R, Adhikari P, Shrestha J, Acharya BP. Response of maize (*Zea mays* L.) hybrids to different levels of nitrogen. Achieves of Agriculture and Environmental Science. 2019;4(3):295-299.
- Eleduma AF, Aderibigbe ATB, Sanni KO, Obabire SO. Effect of cattle manure on soil nutrients dynamics and performances of maize (*Zea mays* L.) grown on sandy loam soil in forest-savannah transition zone southwest Nigeria. International Journal of Horticulture and Food Science. 2020;2(2):05-09.
- Singh C, Singh P, Singh R. Modern Techniques of Raising Field Crops. Oxford and IBH Publishing Company Pvt. Ltd., New Delhi; c2012. p. 84-111.
- Sujatha MG, Lingaraju BS, Palled YB, Ashalatha KV. Importance of integrated nutrient management practices in maize under rainfed condition. Karnataka Journal Agricultural Science. 2008;21(3):334-338.
- Verma NK. Integrated nutrient management in winter maize (*Zea mays* L.) sown at different dates. Journal of Plant Breeding and Crop Science. 2011;3(8):161-167.
- Yadav AK, Chand S, Thenua OVS. Effect of integrated nutrient management on productivity of maize with mungbean intercropping. Global Journal of Bio Science; c2016.
- Yigermal H, Nakachew K, Assefa F. Effect of integrated nutrient application on phenological, vegetative growth and yield-related parameters of maize in Ethiopia. Cogent Food & Agriculture. 2019;5:01-12.
- Raman R, Suganya K. Effect of integrated nutrient management on the growth and yield of hybrid maize. Open Access Journal of Agricultural Research. 2018;3(2):000156.
- Onyia VC, Chime CC, Udeozo PI, Okwesili LC. Concentration levels of organochlorine pesticides (OCPs) residue in soils of selected maize farms within Enugu State, Nigeria